

Bomarea pastazensis (Alstroemeriaceae), an exceptionally small new species from the eastern Andean slopes of Ecuador

John L. Clark¹, Alisson Fierro-Minda^{2,3,4}, Nolan Exe⁵, Mia Johnson⁵, Carrie M. Tribble⁶, Lou Jost⁵

1 Marie Selby Botanical Gardens, 1534 Mound Street, Sarasota, FL 34236, USA

2 Herbario QUSF, Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, USFQ, Quito, Ecuador

3 Herbario QCA, Escuela de Ciencias Biológicas, Pontificia Universidad Católica de Ecuador, Quito, Ecuador

4 Reserva: The Youth Land Trust, Washington DC, USA

5 Reservas de Fundación EcoMinga, Fundación EcoMinga, Baños, Tungurahua, Ecuador

6 School of Life Sciences, University of Hawai'i at Mānoa, Honolulu, HI, USA

Corresponding author: John L. Clark (jclark@selby.org)

Abstract

Recent field research on the eastern slopes of the Andes resulted in the discovery of a new species of *Bomarea* from the Cerro Candelaria Reserve in the Tungurahua province of Ecuador. *Bomarea pastazensis* is the second smallest species in the genus and differs from the smallest by the presence of glutinous trichomes on the ovary, glabrous sepals, and greenish-yellow petals with purple spots. Based on IUCN guidelines, a preliminary conservation status is assigned as Vulnerable (VU).

Resumen

Recientes investigaciones de campo en las estribaciones orientales de los Andes dieron como resultado el descubrimiento de una nueva especie de *Bomarea* en la Reserva Cerro Candelaria en la provincia de Tungurahua en Ecuador. *Bomarea pastazensis* es la segunda especie más pequeña del género y se diferencia por la presencia de tricomas glutinosos en el ovario, sépalos glabros y pétalos verdes con amarillo y manchas moradas. Con base en los lineamientos de la IUCN, se le asigna un estado preliminar de conservación de Vulnerable (VU).

Key words: Alstroemeriaceae, Andes, *Bomarea*, Ecuador, endemism, taxonomy



Academic editor: M. Alejandra Jaramillo

Received: 4 August 2023

Accepted: 10 October 2023

Published: 10 November 2023

Citation: Clark JL, Fierro-Minda

A, Exe N, Johnson M, Tribble CM,

Jost L (2023) *Bomarea pastazensis*

(Alstroemeriaceae), an exceptionally

small new species from the eastern

Andean slopes of Ecuador. *PhytoKeys*

235: 31–42. [https://doi.org/10.3897/](https://doi.org/10.3897/phytokeys.235.110525)

[phytokeys.235.110525](https://doi.org/10.3897/phytokeys.235.110525)

Copyright: © John L. Clark et al.

This is an open access article distributed under terms of the Creative Commons Attribution

License (Attribution 4.0 International – CC BY 4.0).

Introduction

There are more than 100 species of *Bomarea* Mirb. in South America (Hofreiter 2006) and 39 species in Ecuador (Harling and Neuendorf 2003). The description of *Bomarea pastazensis* brings the total diversity to 40 species in Ecuador. The most recent monograph of *Bomarea* was provided by Hofreiter and Tillich (2002) and included an updated classification system based on Baker (1888) that recognized the following four subgenera: *Baccata* Hofr., *Bomarea* Baker, *Sphaerine* (Herb.) Baker, and *Wichuraea* (M. Roemer) Baker. More recently, molecular phylogenetic studies have failed to support the monophyly of these subgenera (Alzate et al. 2008b; Tribble et al. 2022), suggesting that many of the

morphological characters that previously defined infrageneric classification are homoplastic or have evolved convergently. Alzate et al. (2008b) showed that traditionally recognized subgeneric ranks proposed by Baker (1888) and Hofreiter and Tillich (2002) are polyphyletic. Tribble et al. (2022) described three main clades within *Bomarea*, but no updated classification system has yet been proposed. In the present study we recognize the lack of strongly supported subgeneric ranks (Alzate et al. 2008b; Tribble et al. 2022) and discuss the traditional subgenus that most resembles *Bomarea pastazensis*.

Bomarea pastazensis most closely resembles the morphologies associated with the subgenus *Sphaerine* because of the following characters (Hofreiter 2005): erect or hanging habit [mostly non-twining except for *B. coccinea* (Ruiz & Pav.) Baker] (Fig. 1E), resupinate leaves (Figs 1D, 2E), and inferior ovaries (Figs 1A–C, 2A). Recent molecular work suggests that *Sphaerine* is non-monophyletic and some morphological characteristics of this group are likely adaptations to environmental conditions (Alzate et al. 2008b; Tribble et al. 2022). The subgenus *Sphaerine* ranges from the northern Andes of Colombia and Venezuela to Bolivia. In Ecuador, the subgenus is distributed mainly on the eastern Andean slopes. A detailed taxonomic history and overview of *Sphaerine* was provided by Hofreiter (2006), including updated circumscriptions and two new species. The addition of *Bomarea pastazensis* brings the total diversity of subgenus *Sphaerine* to 13 species.

Ecuador is the center of diversity for *Bomarea* (Hofreiter and Tillich 2002; Harling and Neuendorf 2003; Hofreiter 2005; Hofreiter and Rodríguez 2006; Alzate et al. 2008a), which is supported by the presence of 40 of the 120+ currently known species. The other species of *Bomarea* subgenus *Sphaerine* occurring in Ecuador (Hofreiter 2005; Hofreiter and Rodríguez 2006) are: *B. brachysepala* Benth. (Fig. 1A), *B. distichifolia* (Ruiz & Pav.) Baker (Fig. 1B), *B. hieronymi* Pax (Fig. 1), *B. linifolia* (Kunth) Baker, and *B. nervosa* (Herb.) Baker.

The Cerro Candelaria Reserve, owned by the Ecuadorian NGO Fundación EcoMinga, is within the upper Pastaza watershed, an area recently documented for high-levels of local endemism because of the presence of microclimates created by Amazon-Andean airflow currents and the irregular topography (Jost 2004). The Cerro Candelaria Reserve comprises 2800+ hectares, ranging in altitude from 1700 to 3860 m (Reyes-Puig et al. 2013). The private reserve is located within the Llanganates-Sangay corridor, which facilitates a protected corridor for biodiversity between the Llanganates and Sangay National Parks (Fig. 3).

Materials and methods

Plants were photographed and collected during three separate field expeditions to the Cerro Candelaria Reserve by Clark in 2016 (Clark 2016), Exe and Johnson in January 2022, and Exe, Johnson, and Fierro-Minda in November 2022. Specimens were deposited at Pontificia Universidad Católica del Ecuador (QCA), Marie Selby Botanical Gardens (SEL), and Universidad de Guayaquil (GUAY). Digital images were taken of live specimens in the field using an Olympus TG5 and a Nikon D100 DSLR with a Nikon 105 mm lens and a Nikon SB-29s ring flash. Morphological observations and measurements were made from live collections, herbarium specimens, and digital images using the program ImageJ (Schneider et al. 2012).




Figure 1. Some species of Ecuadorian *Bomarea* subgenus *Sphaerine* **A** *Bomarea brachysepala* **B** *Bomarea distichifolia* **C** *Bomarea hieronymi* **D** resupinate leaf (rotated petiole indicated with white arrow) of *Bomarea hieronymi* **E** terrestrial habit (non-twining) of *Bomarea hieronymi*. Photos **A**, **B** by N. Exe, **C–E** by J.L. Clark (J.L. Clark 17350).

We assessed the extinction risk of *Bomarea pastazensis* following the IUCN Red List Categories and Criteria (IUCN 2012) and guidelines of the IUCN Standards and Petitions Committee (2022). We considered observations, collection localities, and population estimate from fieldwork. Species extent of occurrence (EEO) and area of occupancy (AOO) were calculated using GeoCAT (Bachman et al. 2011) with the default setting of 2 km² grid.

A note on terminology

Some botanical terminology has been used inconsistently in previous descriptions of *Bomarea*, and other terminology is specific to *Bomarea*. Here, we clarify our use of potentially confusing terms to ensure that our definitions are unambiguous while linking the description to other literature. Botanical glossaries usually refer to sepals as the outer whorl of the perianth and petals as the inner whorl of the perianth (e.g., Harris and Harris 2006; Pell and Angell 2016). In contrast, ‘tepal’ is used when describing cases where the outer and inner perianth whorls are identical or when perianth whorls are not readily differentiated. Some *Bomarea* taxonomists use the terms inner and outer tepals in their descriptions of the perianth (Hofreiter 2005; 2006; Hofreiter and Rodríguez 2006) and others use the terms sepals and petals (Harling and Neuendorf 2003). The description of *Bomarea pastazensis* uses sepals and petals to provide greater specificity because the two perianth layers are easily differentiated. Furthermore, we use the terms adaxial and abaxial following the definition of these terms to refer to the portion of the leaf facing towards or away from, respectively, the structure-bearing organ (the stem) during development (Eckel 2011). When referring to the surface of resupinate leaves (rotating 180 degrees) of *Bomarea pastazensis*, abaxial is the upper surface of the leaf, and adaxial is the lower surface of the leaf, terminology that is consistent with other *Bomarea* descriptions (Hofreiter and Lyshede 2006). Following Hofreiter (2005), we use the term “claw” to refer to the horn-like thickened sepal apex in many *Bomarea* species.

Taxonomic treatment

***Bomarea pastazensis* J.L.Clark, Fierro-Minda & N.Exe, sp. nov.**

urn:lsid:ipni.org:names:77330582-1

Figs 2, 4

Diagnosis. Similar to *Bomarea pumila* Griseb. ex Baker, differing in plant height reaching 10–14 cm (vs. 5–8 cm high in *B. pumila*), pubescent stem (vs. glabrous stem in *B. pumila*), the presence of ciliate leaf margin (vs. glabrous leaf margin in *B. pumila*), and the presence of dark spots on the petals (vs. no spotting on the petals in *B. pumila*).

Type. ECUADOR. Tungurahua: Cantón Baños, Parroquia Río Verde, Cerro Candelaria Reserve (Fundación EcoMinga), upper Pastaza watershed, 1°28'39.33"S, 78°17'53.61"W, 3642 m, 10 Mar 2016, J.L. Clark 14839 (holotype: QCA! [245371]; isotype: SEL! [079072]).

Description. Terrestrial or epiphytic herb. **Rhizome** short with multiple underground fusiform to globose root tubers, pale and heavily stippled with dull reddish-purple spots, 5–9 × 3–4 mm, surface striate, yellowish-brown with red spots




Figure 2. *Bomarea pastazensis* **A** lateral view of flower **B** front view of flower featuring mature androecium **C** mature gynoecium **D** petal **E** adaxial leaf surface **F** mature flowers with foliage **G** habitat **H** tuberous roots. Photos **A** by J.L. Clark (J.L. Clark 14839), **B–H** by N. Exe (N. Exe et al. 2).




Figure 3. Distribution map of *Bomarea pastazensis*. Red dots indicate collections or observations.

(Fig. 2H). **Stem** erect, 10–14 cm long, ca. 0.13 cm in diameter, slender, terete, base pubescent, apex puberulous, internodes 0.6 cm long. **Leaves** alternate; blade ovate, 1.4–1.8 × 0.4–0.7 cm, base rounded to obtuse, apex acute, abaxially glabrous, adaxially light green, suffused with whitish-translucent trichomes, ca., 0.2 mm long, clustered along veins; blades with 5–7 prominent parallel veins, raised below and slightly raised above; margin hyaline, slightly revolute with unicellular trichomes to 0.2 mm long; petioles resupinate and canaliculate, 0.1–0.4 cm long, basal leaves reduced to scales. **Flowers** produced from the stem apex, usually one (rarely two) apical flower(s) per stem.; pedicels 1–1.4 cm long, terete, with brownish red to dark purple trichomes, bisexual, actinomorphic, epigynous. **Sepals** 3, 1.0–1.3 × 1.0–1.2 cm, each sepal with 6 parallel veins, broadly ovate, apically rounded, bright red to orange-red with a 1 mm green to black claw, inner and outer surfaces glabrous. **Petals** 3, basally constricted and caniculate, distally obovate to broadly spatulate, 1.5–1.6 × 0.6–0.7 cm, greenish yellow, puberulous at base, inner surface with 3 large reddish-brown spots, outer surface uniformly yellow (occasionally yellow suffused with red from the inner spots). **Androecium** of 6 free stamens, 0.8–0.9 cm long, thickened near center of filaments; anthers pseudo-basifixed, 0.3 × 0.1 cm, fusiform; pollen grains lilac. **Gynoecium** comprised of three fused carpels, ovary 0.5 × 0.4 cm with surface covered with glutinous trichomes, style ca. 1 cm long, stigma with three circinate lobes. **Fruits** not observed.

Ecology. Found growing on *Sphagnum* and mossy cushions at ground level and epiphytically (up to 2.5 m high) in high elevation cloud forest and paramo



Figure 4. Holotype of *Bomarea pastazensis*, J.L. Clark 14839 (QCA).

(observed from 3235 to 3700 m). Plants growing in partially shaded areas to full sunlight. Stems erect to hanging, with flowers often found on or slightly above ground level. Flowers protandrous, commonly with one flower per stem but occasionally two.

Phenology. Observed in flower in January, March, July, and November. Approximately 20 individual plants with mature flowers were located during an expedition in November of 2022 and fewer individuals with mature flowers were observed in January of 2022. Fruits not documented.

Etymology. The specific epithet, *pastazensis*, reflects the watershed of the type locality that includes Río Pastaza and adjacent tributaries.

Distribution and preliminary assessment of conservation status. *Bomarea pastazensis* is endemic to the upper Pastaza watershed, located in the eastern Andean slopes of Ecuador. The first documented population was inside Cerro Candelaria Reserve in Tungurahua province (Fig. 3). The Cerro Candelaria Reserve is a private reserve of 2800+ hectares managed by Fundación EcoMinga. It is bordered on the north by EcoMinga's Naturetrek Reserve, and the south by Parque Nacional Sangay. Along with Fundación EcoMinga's Machay and Naturetrek Reserves to the north, it forms a protected corridor between Parque Nacional Sangay and Parque Nacional Llanganates. Cerro Candelaria Reserve was founded by Fundación EcoMinga in 2007 and financed by the World Land Trust (UK); the majority of the reserve is undisturbed by anthropogenic activity. A diverse flora and fauna has been recorded here, including many endemic species and species of high conservation priority (Jost 2004; Reyes-Puig et al. 2013), such as *Blakea attenboroughii* Penneys & L.Jost (Melastomataceae) which is endemic to the lower elevation forests of Bosque Protector Cerro Candelaria. Several recently described species in the Orchidaceae share a similar geographic distribution to *Bomarea pastazensis* (Fig. 3), include *Teagueia barbeliana* L.Jost & Shepard, *T. puroana* L.Jost & Shepard, *T. kostoglouana* L.Jost & Shepard, *T. lizziefinchiana* L.Jost & Shepard, and *T. anitana* L.Jost & Shepard (Jost and Shepard 2011; 2017). Additional surveys in Cerro Candelaria and in the surrounding upper Rio Pastaza watershed are likely to yield many more undescribed species, and potentially additional populations of *Bomarea pastazensis*. The distribution of this species highlights the importance of the Llanganates-Sangay corridor for allowing species gene flow and connectivity among populations (Ríos Alvear and Reyes-Puig 2015). A second documented population of *Bomarea pastazensis* was provided by digital images from EcoMinga's park guard Eduardo Peña (July 2023) in the Llanganates National Park, north of Río Pastaza (Fig. 3). We calculated the AOO=12 km² from Peña's observation and the type collection (Fig. 3). Based on the available information and according to the IUCN Red List criteria (IUCN 2012; IUCN Standards and Petitions Committee 2022), *Bomarea pastazensis* is preliminarily assessed as Vulnerable (VU) based on a limited area of occupancy (IUCN criterion D2 where AOO <20 km²) and dependence on conservation efforts for its continued survival. The only documented populations of *Bomarea pastazensis* are located inside protected areas, at elevations that are used for agriculture in unprotected parts of the Pastaza province. Effective conservation of this and the other unique species of the Pastaza watershed will require constant vigilance.

Comments. *Bomarea pastazensis* differs from other congeners by a distinctive pattern of three dark spots on the petals (Fig. 2D), widely opened flowers,




Figure 5. *Bomarea pumila* A–D field images from Peru E syntype (K! [K000400912]) of *Bomarea pumila* of W. Lechler 2240 from Sachapata, Peru. Photos A–D from Lucely L. Vilca Bustamente E from The Herbarium Catalogue, Royal Botanic Gardens, Kew. The Board of Trustees of the RBG, Kew.

Table 1. General geographic distribution (names in parentheses indicate Ecuadorian province) and comparison of morphological characters between *Bomarea pastazensis* and *B. pumila*.

	<i>Bomarea pumila</i>	<i>Bomarea pastazensis</i>
Habit	erect	erect to hanging
Plant height	5–9 cm long	10–14 cm long
Stem surface	glabrous	pubescent
Leaf – relative size	leaves not uniform on stem; central leaves relatively larger compared with basal and apical leaves	leaves relatively uniform on stem
Leaf margin	glabrous	ciliate
Pedicle length	1.5–4 cm long	1–1.4 cm long
Flower length	0.8–1.5 cm long	1.5–1.6 cm long
Flower width during anthesis	2–3 cm wide	1.5–2 cm wide
Sepal surface	conspicuously pubescent	sparsely pubescent at base only
Sepal horn	white to reddish horn at apex	green to black horn at apex
Petal coloration (=inner tepals)	yellow with a red stripe and green tip.	green suffused with yellow with three prominent red spots at the base
Distribution	Peru and Bolivia	Ecuador (Tungurahua)

and small size. *Bomarea pumila* and *B. pastazensis* share similar traits such as their small size in comparison with the rest of *Bomarea* species, the presence of pubescence throughout their above-surface organs, and the color patterns of the flowers (Figs 2, 5; Table 1). *B. pastazensis* is slightly bigger in size (10–14 cm tall) than *B. pumila* (5–9 cm tall). In contrast, *B. pastazensis* is sparsely pubescent at the base of the sepals, while the sepals of *B. pumila* are uniformly densely pubescent (Fig. 5). The petals of *B. pumila* are not spotted, which differentiates it from *B. pastazensis* which has a three-spotted pattern on the petals (Fig. 2B). The two species are geographically isolated with *B. pastazensis* endemic to the Pastaza watershed on the eastern Andean slopes in central Ecuador (Tungurahua province) and *B. pumila* from Central Peru to northern Bolivia. Table 1 provides a summary of distribution and useful characters for differentiating *B. pastazensis* and *B. pumila*.

Additional specimens examined. ECUADOR. Tungurahua: Cantón Baños, Parroquia Río Verde, Bosque Protector Cerro Candelaria (Fundación EcoMinga), upper Pastaza watershed, Cerro Candelaria summit trail, just below paramo, 1°28'39.33"S, 78°17'53.61"W, 3150–3827 m, 1 Jan 2022, N. Exe, M. Johnson & A. Fierro-Minda 2 (GUAY).

Acknowledgements

We thank the Ministry of the Environment of Ecuador for providing permits for specimen collection and transportation (Research permit Aves y Conservación N° 007-2018-IC-FLO-FAU and mobilization permit Aves y Conservación 005-FLO-2019-DPAP-MA). We thank Lucely L. Vilca Bustameante (Instituto Nacional de Investigación en Glaciares y Ecosistemas de Montaña, Cusco, Peru) for sharing her field images of *Bomarea pumila* (Fig. 5). We are grateful to Guillermo Peña for assisting with the expedition and collection of specimens, as well as providing

supplemental locality information. We thank the Science Collections Digitisation Project at the Royal Botanic Gardens, Kew for facilitating an image of the syntype for *Bomarea pumila* (Fig. 5). We are grateful to Alan W. Meerow and an anonymous reviewer for providing helpful feedback on an earlier version of the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

No funding was reported.

Author contributions

John Clark made the type collection in 2016. Nolan Exe, Mia Johnson, and Alisson Fierro-Minda conducted additional fieldwork in 2022. All authors contributed to writing the manuscript. Jost wrote the IUCN assessment. Tribble facilitated the use of terminology specific to Alstroemeriaceae.

Author ORCIDs

John L. Clark  <https://orcid.org/0000-0002-1414-6380>

Alisson Fierro-Minda  <https://orcid.org/0000-0001-7169-2989>

Nolan Exe  <https://orcid.org/0000-0001-8847-7978>

Mia Johnson  <https://orcid.org/0000-0002-4822-0606>

Carrie M. Tribble  <https://orcid.org/0000-0001-7263-7885>

Lou Jost  <https://orcid.org/0000-0002-5850-0716>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Alzate F, Quijano-Abril MA, Morrone JJ (2008a) Panbiogeographical analysis of the genus *Bomarea* (Alstroemeriaceae). *Journal of Biogeography* 35(7): 1250–1257. <https://doi.org/10.1111/j.1365-2699.2008.01896.x>
- Alzate F, Mort ME, Ramirez M (2008b) Phylogenetic analyses of *Bomarea* (Alstroemeriaceae) based on combined analyses of nrDNA ITS, *psbA-trnH*, *rpoB-trnC* and *matK* sequences. *Taxon* 57(3): 853–862. <https://doi.org/10.1002/tax.573014>
- Bachman S, Moat SJ, Hill A, de la Torre J, Scott B (2011) Supporting Red List threat assessments with GeoCAT: Geospatial conservation assessment tool. *ZooKeys* 150: 117–126. <https://doi.org/10.3897/zookeys.150.2109>
- Baker JG (1888) *Handbook of the Amaryllideae*. George Bell & Sons, London, UK.
- Clark JL (2016) The 2016 Lawrenceville School in Ecuador Program. *Gesneriads* 66: 13–18.
- Eckel PM (2011) *A Grammatical Dictionary of Botanical Latin*. Missouri Botanical Garden. <http://www.mobot.org/mobot/LatinDict/search.aspx>

- Harling G, Neuendorf M (2003) Alstroemeriaceae. 71. In: Harling GW, Andersson L (Eds) Flora of Ecuador. University of Göteborg, Sweden, 3–108.
- Harris JG, Harris MW (2006) Plant identification terminology: an illustrated glossary. Spring Lake Publishing, Spring Lake, Utah, USA, 206 pp.
- Hofreiter A (2005) The genus *Bomarea* (Alstroemeriaceae) in Bolivia and Southern South America. Harvard Papers in Botany 9: 343–374.
- Hofreiter A (2006) Revision of *Bomarea* subgenus *Sphaerine* (Alstroemeriaceae). Nordic Journal of Botany 24(2): 117–141. <https://doi.org/10.1111/j.1756-1051.2004.tb00827.x>
- Hofreiter A, Lyshede OB (2006) Functional leaf anatomy of *Bomarea* Mirb. (Alstroemeriaceae). Botanical Journal of the Linnean Society 152(1): 73–90. <https://doi.org/10.1111/j.1095-8339.2006.00540.x>
- Hofreiter A, Rodríguez EF (2006) Alstroemeriaceae en Perú y áreas vecinas. Revista Peruana de Biología 13(1): 5–69. <https://doi.org/10.15381/rpb.v13i1.1765>
- Hofreiter A, Tillich H-J (2002) The delimitation, infrageneric subdivision, ecology, and distribution of *Bomarea* Mirbel. (Alstroemeriaceae). Feddes Repertorium 113(7–8): 528–544. <https://doi.org/10.1002/fedr.200290005>
- IUCN (2012) IUCN Red List Categories and Criteria: Version 3.1. 2nd edn. IUCN, Gland, Switzerland and Cambridge.
- IUCN Standards and Petitions Committee (2022) Guidelines for using the IUCN Red List Categories and Criteria. Version 15.1. <https://www.iucnredlist.org/resources/redlist-guidelines>
- Jost L (2004) Explosive local radiation of the genus *Teagueia* (Orchidaceae) in the Upper Pastaza Watershed of Ecuador. Lyonia 7: 41–47.
- Jost L, Shepard A (2011) Two new species of *Teagueia* (Orchidaceae: Pleurothallidinae) from East-Central Ecuador. Lankesteriana 1(11): 9–14. <https://doi.org/10.15517/lank.v11i1.18310>
- Jost L, Shepard A (2017) Four new *Teagueia* (Orchidaceae: Pleurothallidinae) from the Upper Río Pastaza watershed of East-Central Ecuador. Lankesteriana 2(17): 261–268. <https://doi.org/10.15517/lank.v17i2.30159>
- Pell SK, Angell B (2016) A botanist's vocabulary: 1400 terms explained and illustrated. Timber Press, Portland, Oregon, USA, 224 pp.
- Reyes-Puig J, Ramírez S, Yáñez-Muñoz M, Morales M, Recalde L (2013) Bosque Protector Cerro Candelaria: Una isla de biodiversidad entre las nubes. In: MECN, JOCOTOCO y ECOMINGA. Herpetofauna en Áreas prioritarias para la conservación: El sistema de Reservas Jocotoco y EcoMinga. Monografía 6. Serie de publicaciones del Museo Ecuatoriano de Ciencias Naturales (MECN), Fundación para la Conservación Jocotoco, and Fundación EcoMinga. Quito, Ecuador, 196–211.
- Ríos Alvear G, Reyes-Puig C (2015) Corredor ecológico Llanganates-Sangay: Un acercamiento hacia su manejo y funcionalidad. Yachana Revista Científica 4: 11–21.
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nature Methods 9(7): 671–675. <https://doi.org/10.1038/nmeth.2089>
- Tribble CM, Alzate-Guarín F, Gándara E, Vartoumian A, Burleigh JG, Zenil-Ferguson R, Specht CD, Rothfels CJ (2022) The rapid radiation of *Bomarea* (Alstroemeriaceae: Liliales), driven by the rise of the Andes. BioRxiv, 2022–09. <https://doi.org/10.1101/2022.09.15.507859>