

Gentiana mopanshanensis (Gentianaceae), a new species from Yunnan, southwest China

Tao Chen¹ , Ting-Ting Wang¹ , Shao-Yun Liu¹ , Huan-Chong Wang^{1,2} 

1 School of Ecology and Environmental Science, Yunnan University, Kunming 650500, Yunnan, China

2 Herbarium of Yunnan University, Kunming 650091, Yunnan, China

Corresponding author: Huan-Chong Wang (hchwang@ynu.edu.cn)

Abstract

Gentiana mopanshanensis, a new species of the family Gentianaceae is here described and illustrated. This species is presently known only from the Mopanshan Mountain, Yunnan Province, southwest China. Phylogenetic analysis based on ITS sequence data has shown that this new species is a member of the series *Fimbriatae* of the section *Chondrophyliae*. Morphologically, it mostly resembles *G. mairei* and *G. panthaica*, but differs clearly from the latter two species in the shape and size of the leaves, and the characters of the corolla throat and plicae.

Key words: Diffuse herbs, endemism, ITS sequence, Mopanshan Mountain, series *Fimbriatae*, Yunnan-Guizhou Plateau

Introduction



Academic editor: Lorenzo Peruzzi

Received: 29 January 2024

Accepted: 6 March 2024

Published: 20 March 2024

Citation: Chen T, Wang T-T, Liu S-Y, Wang H-C (2024) *Gentiana mopanshanensis* (Gentianaceae), a new species from Yunnan, southwest China. PhytoKeys 239: 215–228. <https://doi.org/10.3897/phytokeys.239.119800>

Copyright: © Tao Chen et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](#)).

The genus *Gentiana* L. belongs to the family Gentianaceae and comprises more than 360 species that are distributed worldwide in the temperate and alpine regions, especially in Europe, Asia and North America (Ho and Liu 1990; Yuan et al. 1996; Yuan and Küpfer 1997). The plants of this genus are typical mountain plants, and most of the species are adapted to alpine habitats (Ho et al. 1996). The greatest diversity of *Gentiana* is found in the Tibeto-Himalayan region, where more than 250 species are native (Ho and Liu 2001). Relevant biogeographic studies have identified this region as the source area for *Gentiana* and related genera (Favre et al. 2016; Matuszak et al. 2016). The phylogenetic relationships within the genus *Gentiana* are now still incompletely clear, and its classification remains controversial (Favre et al. 2014, 2020; Sun et al. 2018; Sun and Fu 2019; Fu et al. 2021, 2022). Many *Gentiana* species have ornamental value and are also of pharmaceutical interest due to their interesting phytochemical properties (Mariana et al. 2013; Mirzaee et al. 2017; Jiang et al. 2021).

China is endowed with numerous species of *Gentiana*, including many endemics (Ho and Pringle 1995). While members of *Gentiana* can be found throughout the country, they are mostly concentrated in the Southwest Mountainous area, which includes the Qinghai-Tibet Plateau and Hengduan Mountains (Ho et al. 1988; Bai 2000). Ho and Pringle (1995) recognized 248 species

from China in their "Flora of China", which accounts for more than 2/3 of the genus' total species. Nevertheless, new species have been frequently discovered in China in recent years (e.g., Hsieh et al. 2007; Yang et al. 2008; Ho and Liu 2010; Wu et al. 2012; Yu et al. 2012; Yang et al. 2020; Favre et al. 2022). These new discoveries highlighted the need for continued field exploration and taxonomic research in this area.

During recent field surveys in Mopanshan Mountain, Xinping County, Yunnan Province, southwest China, we discovered an unknown species of *Gentiana*. After a phylogenetic analysis using ITS sequences to infer its systematic position and a detailed comparison with morphologically similar species, it became clear that this plant represents a distinct new species.

Materials and methods

Morphological analyses

This study including plant collection, specimen preparation, observation, and analysis followed the normal practice of plant taxonomic survey and herbarium taxonomy (Davis and Heywood 1963). Morphology of the new species was studied based on observation of living plants and specimens housed at YUKU. Digital images of type specimens of genus *Gentiana* are available at the JSTOR Global Plants (<http://plants.jstor.org/>) and at the Chinese Virtual Herbarium (<http://www.cvh.ac.cn/>); in addition, the collections housed at GBIF, KUN and YUKU were examined and compared with the new species. The dried specimens were examined under stereomicroscopes for morphological studies, and various organs were measured using rulers and metric vernier calipers. Terminology followed Ho and Liu (2001), Beentje (2010) and Mayfield (2021).

Phylogenetic study

To determine the phylogenetic position of the putative new species, the internal transcribed spacer region (ITS) of the nuclear ribosomal DNA was used as the molecular marker. The total genomic DNA of this new species is extracted from silica-gel dried leaves using the DNA secure plant kit (Tiangen, Amsterdam, Netherlands). The PCR protocol followed by Shabir et al. (2022). The ITS primers used in this study were ITS4 and ITS5, as described by Yuan and Küpfer (1995) and He et al. (2016). The PCR products were bidirectionally sequenced with the same primers used for PCR amplifications in an ABI 3730 XL DNA Analyzer (Applied Biosystems) at the Kunming Branch of Beijing Qingke Biotechnology Co., Ltd. (Yunnan, China).

There are a total of 44 species of *Gentiana* to be used, which are representatives of most sections of *Gentiana*, including *G. sect. Chondrophyliae* Bunge, *G. sect. Frigida* Kusnezow, *G. sect. Monopodiae* (H. Smith) T. N. Ho, *G. sect. Phyllocalyx* T. N. Ho and *G. sect. Stenogyne* Franchet. In addition, *Swertia rosulata* (Baker) Klack, *Halenia taruga-gasso* Gilg and *Gentianella gentianoides* (Franchet) H. Smith were selected as outgroups. The dataset for phylogenetic analysis consists of 60 taxa, 46 of which were obtained from the GenBank. Voucher specimen and GenBank accession information for taxon are listed in Appendix 1.

All sequences were aligned with MAFFT (Katoh and Standley 2013) using ‘auto’ strategy and normal alignment mode. Gap sites were removed with trimAI (Capella-Gutiérrez et al. 2009) using the “-automate” command. The best-fitting substitution models SYM+I+G model for Bayesian inference were selected using ModelFinder (Kalyaanamoorthy et al. 2017) in BIC criterion. MrBayes 3.2.6 (Ronquist et al. 2012) was used to conduct Bayesian phylogenetic analyses. Runs were performed for 5 million generations with a sampling of trees every 500 generations. The initial 25% of sampled data were discarded as burn-in.

Results and taxonomic treatment

Gentiana mopanshanensis Huan C. Wang & Tao Chen bis, sp. nov.

urn:lsid:ipni.org:names77338758-1

Figs 1–3

Type. CHINA. Yunnan Province: Xinping County, Mopanshan Mountain, near the top of mountain, alt. 2480 m, 23°56'23"N, 101°59'23"E, 3 April 2023, in flower, H. C. Wang et al. XP19775 (Holotype: YUKU!; isotypes: YUKU!).

Diagnosis. *Gentiana mopanshanensis* is distinguishable from all other similar species of the genus by the combination of its rosulate basal leaves lanceolate to gladiate, up to 5 (6) cm long, caudine leaves lanceolate or linear-lanceolate, throat of corolla blue maculate, plicae with 5–10 fimbriations, and fimbriation irregular in length, usually 0.5–2 mm long.

Etymology. The specific epithet “*mopanshanensis*” is derived from the type locality of the new species, the Mopanshan Mountain, and the Latin suffix *-ensis*, indicating the place of origin or growth.

Description. Biennial herbs, diffuse, 5–15 cm in height. Root slightly fleshy, 4–6 cm long, with conspicuous rootstock. Stems yellow-green or purplish-red, smooth, much branched at base; branches procumbent or ascending. Basal leaves rosulate, persistent at anthesis, sessile or subsessile; blades lanceolate to gladiate, (1–) 3–5 (–6) cm long, 0.2–0.7 (–1) cm wide, both surfaces densely and minutely papillate, apex acuminate, margin transparent, densely denticulate, basal veins 1–3, distinct, midvein convex beneath. Cauline leaves opposite, semiamplexicaul, base proximally compounded, petiole tube 1–1.5 mm long; blades lanceolate or linear-lanceolate, 0.3–2 cm long, 0.1–0.6 cm wide, apex acuminate, margin transparent, densely denticulate, both surfaces densely and minutely papillate, basal veins 1–3, midvein convex beneath. Flowers solitary, terminal on branch. Pedicels yellowish green, glabrous, 3–12 mm long, purplish-red striped. Calyx 5–8 mm long, obconic, yellow-green; tube campanulate, 4–5 mm long, 2–3 mm in diameter, slightly longer than lobes; lobes 5, acicular or subulate, 1.5–3 mm long, papillate on margin; veins ridged on abaxial surface, decurrent towards calyx tube; sinus between lobes obtuse to sub-rounded. Corolla 8–14 mm long, 5–8 mm in diameter, trumpet-shaped, blue-white, outside with copper-green stripes, inside with deeply blue spots in throat; tube tubular, 5–8 mm long, 4–6 mm in diameter, golden inside; lobes ovate to broadly ovate, 2–3 mm long, 2–3 mm wide, apex acute, margin entire; plicae triangular-ovate, 2–2.5 mm long, apex irregularly laciniate, fimbriate, with 5–10 fimbriations, usually 0.5–2 mm long. Stamens 5, filaments filiform-subulate, 3–6 mm long, inflated at middle, inserted in lower middle of corolla tube; anthers rectan-




Figure 1. *Gentiana mopanshanensis* sp. nov. (Drawn by Ting T. Wang) A habit B flower (front view) C dissected corolla D dissected calyx E seed.

gular-rounded, 0.6–1 mm long. Ovary stipitate, ellipsoid or fusiform, 2.5–3 mm long, apex obtuse, base attenuate; style clavate, 0.5–1 mm long, stigma bifid, extrorse. Capsules obovate, exerted beyond persistent corolla, gynophore up to 15 mm long, narrowly winged on both margins. Seeds ovate-triangular, up to 1.5 mm long, yellowish-brown, densely striato-reticulate on seed coats.

Phenology. *Gentiana mopanshanensis* has a long flowering and fruiting period. It starts flowering in March and continues until August, and its fruiting period is from May to September.

Distribution and habitat. *Gentiana mopanshanensis* is currently only found in the Mopanshan Mountain (Fig. 4), which is located in the southeast of Xinping County, at the southwestern edge of the Yunnan-Guizhou Plateau. The mountain is situated east of the Yuanjiang River valley and has an elevation




Figure 2. *Gentiana mopanshanensis* sp. nov. (Photographed by HCW and TC) **A** habitat **B** habit **C** root **D** leaf blade (side view, showing adaxially densely and minutely papillate and transparent denticulate on margin) **E** closed flower **F** flower (front view) **G** dissected flower (showing the stamens and style) **H** mature fruit protruding from persistent corolla **I** stamens **J** pistils **K** seed.



Figure 3. Holotype of *Gentiana mopanshanensis* sp. nov. (YUKU-05008414).




Figure 4. Geographical distribution of *Gentiana mopanshanensis* sp. nov. (red dot).

ranging from 1370 to 2611 meters. *G. mopanshanensis* usually occurs at elevations between 2400 and 2550 meters and mainly grows in wet meadows near the peak of the mountain. It can also be occasionally found under the thickets predominated by *Lithocarpus variolosus* Chun (Fagaceae) and *Quercus guyavifolia* H.Lév. (Fagaceae). In meadow habitats, this new species is commonly associated with *G. praticola* Franchet (Gentianaceae), *Polygala dunniana* H.Lév. (Polygalaceae), *Arundinella hookeri* Munro ex Keng (Gramineae), *Fragaria nilgerrensis* Schlecht. ex J. Gay (Rosaceae), *Roscoea tibetica* Batalin (Zingiberaceae) and *Bistorta paleacea* Yonek. et H.Ohashi (Polygonaceae).

Molecular phylogenetics. The ITS sequence region of *Gentiana mopanshanensis* comprises 625 base pairs with a GC content of 57.12%. The alignment of 60 ITS sequences resulted in a matrix of 662 total characters, of which 344 are constant, 93 of the variable characters are singleton sites and 225 characters are parsimony informative sites.

As shown in the phylogenetic tree (Fig. 5), phylogenetic analyses using the ITS sequence data demonstrated that the new species belongs to a clade representing the *Gentiana* section *Chondrophyliae* with maximum support. In this clade, *G. mopanshanensis* falls within the subclade corresponding to series *Fimbriata* Marq. with 0.993 posterior probabilities. It constituted a monophyletic lineage with *G. panthaica* Prain et Burkhill and *G. mairei* H.Lév. with maximum support (PP = 1) and were resolved as sister to them. The phylogenetic result is also supported by the morphological characteristics.




Figure 5. Mrbayes tree of *Gentiana* based on ITS sequences showing phylogenetic placement of *G. mopanshanensis*. Mrbayes posterior probabilities are shown near the nodes. *G. mopanshanensis* is marked by a red box.

Discussion. Based on phylogenetic analyses, *Gentiana mopanshanensis* should be assigned to the series *Fimbriatae* of the section *Chondrophyliae*. Its placement within this series is also supported by its particular morphological characters: stems much branched at the base, basal leaves well developed, leaf blades and calyx with densely and minutely papillate, calyx lobes acicular or subulate, plicae apex fimbriate, obovate capsule with strong and broad wings at apex.

This new species is most similar to *Gentiana mairei* and *G. panthaica* in terms of habit, and flower shape and size, especially plicae apex fimbriate. However, *G. mopanshanensis* can be distinguished from *G. mairei* by its biennial (vs. annual) habit, basal leaves lanceolate to gladiate (vs. ovate to ovate-elliptic), (1–) 3–5 (–6) cm (vs. 0.7–1.4 cm) long, 0.2–1 cm (vs. 0.3–0.7 cm) wide, caudine leaves lanceolate or linear-lanceolate (vs. ovate-triangular to ovate-lanceolate), with blue spots (vs. blackish, white or pale yellow stripes) in throat, plicae with 5–10 (vs. 8–10) fimbriations, irregular (vs. nearly equal) in length, usually 0.5–2 mm (vs. 2–2.5 mm) long. *G. mopanshanensis* differs from *G. panthaica* in biennial (vs. annual) habit, basal leaves lanceolate to gladiate (vs. ovate-elliptic to ovate), caudine leaves lanceolate or linear-lanceolate (vs. lanceolate, narrowly elliptic or ovate-triangular), plicae with 5–10 (vs. 15–20) fimbriations, filament laciniate (vs. filiform, curly). A detailed morphological comparison between these three species is summarized in Table 1.

Additional specimens examined. *Gentiana mopanshanensis*. China. Yunnan: Xinping County, Mopan Mountain, alt. 2509 m, 13 August 2012, in flower and

Table 1. Morphological comparison of *G. mopanshanensis*, *G. mairei*, *G. panthaica*.

Characters	<i>G. mopanshanensis</i>	<i>G. mairei</i>	<i>G. panthaica</i>
Habit	biennial	annual	annual
Leaf			
-Basal leaf	lanceolate to gladiate 3–5 (6) cm × 0.2–0.7 (1) cm	ovate to ovate-elliptic, 7–1.5 cm × 3–0.7 cm	ovate-elliptic to ovate, 0.9–2 cm × 0.4–1 cm
-Caudine leaf	lanceolate or linear-lanceolate, 0.3–1.5 cm × 0.1–0.4 cm	ovate-triangular to ovate-lanceolate 0.55–1.1 cm × 0.25–0.4 cm	lanceolate, narrowly elliptic or ovate-triangular, 0.6–0.8 cm × 0.25–0.3 cm
Flower			
-Calyx			
-Tube	campanulate, 4–5 mm long	obconic, 3–4 mm long	obconic, 3–4 mm long
-Lobe	acicular or subulate, 1.5–3 mm long	filiform-subulate, 2–2.5 mm long	filiform to filiform-conical, 1.5–3 mm long
-Coroll	trumpet-shaped, 8–14 mm long, 5–8 mm in diam.	obconic 8.5–12 mm long, 5–8 mm in diam.	obconic 8–14 mm long, 5–10 mm in diam.
-Lobe	ovate to broadly ovate, 2–3 mm long	ovate-orbicular, 2.5–3 mm long	ovate, 2.5–3.5 mm long
-Plicae	triangular-ovate, with 5–10 fimbriae	ovate-oblong, with 8–10 fimbriae	ovate, with 15–20 fimbriae
-Stamen			
-Filament	filiform-subulate, 3–6 mm long	filiform, 3–3.5 mm long	filiform-subulate, 3–4 mm long
-Anther	rectangular-rounded	ellipsoid	ellipsoid
Pistil			
-Ovary	ellipsoid or fusiform, 2.5–3 mm long	ellipsoid, 2.5–3 mm long	ellipsoid, 3–4 mm long
-Style	clavate, 0.5–1 mm long	linear, 0.7–1.5 mm long	clavate, 0.7–1.2 mm long
Fruit	obvoid, 4–7 mm long	obvoid, 4.5–5.5 mm long	obvoid, 4–5 mm long
Seed	ovate-triangular, 1–1.5 mm long	ellipsoid, 1.3–1.5 mm long	ellipsoid, 1.3–1.5 mm long

fruit, *Xinping County Census team 5304270757* (IMDY0019083); same location, alt. 2406 m, 18 June 2023, in flower and fruit, *T. Chen et al. XP23338* (YUKU).

Gentiana mairei. China. Yunnan: Dali City, Cangshan Mountain, alt. 3800 m, 22 July 2009, *Z.J.Yin et al. 1631* (KUN-1220364); Jingdong County, Wuliangshan Mountain, alt. 3100 m, 19 November 1956, *B.Y.Qiu 53823* (KUN-00088281).

Gentiana panthaica. China. Yunnan: Heqing County, Mae Shan, 15 August 2020, *Q.P. Wang et al. HQ 8930* (YUKU); Nanjian County, Wuliangshan Mountain, alt. 2270 m, 24 March 2012, *E.D. Liu et al. 3587* (KUN-1224606); Dali City, Cangshan Mountain, alt. 3800 m, 15 July 2009, *Z.J.Yin et al. 1362* (KUN-1220362); same location, 13 July 2009, *Z.J.Yin et al. 1111* (KUN-1220361); Luquan County, Daheiqing, alt. 3150 m, 2 July 1990, *R.F.Fang et al. 83* (KUN-551847).

Acknowledgements

We are grateful to the editors of PhytoKeys and the anonymous reviewers for critical comments on the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by grants-in-aid from the National Natural Science Foundation of China (NSFC) (Grant No.31960040) and the Second Tibetan Plateau Scientific Expedition and Research (STEP) program (2019QZKK0502).

Author contributions

Investigation: HCW, TC, SYL, TTW. Writing - original draft: TC. Writing - review and editing: HCW, TC.

Author ORCIDs

Tao Chen  <https://orcid.org/0009-0004-1045-1662>

Ting-Ting Wang  <https://orcid.org/0009-0000-1593-1738>

Shao-Yun Liu  <https://orcid.org/0009-0001-6524-7632>

Huan-Chong Wang  <https://orcid.org/0000-0001-8562-8849>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bai PY (2000) *Gentiana* L. In: Wu ZY (Ed.) *Flora Yunnanica*, Vol. 11. Science Press, Beijing, 47–634.
- Beentje H (2010) *The Kew Plant Glossary: An Illustrated Dictionary of Plant Terms*. Royal Botanic Gardens, Kew, 160 pp.

- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics (Oxford, England) 25(15): 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
- Davis PH, Heywood VH (1963) Principles of Angiosperm Taxonomy. Oliver and Boyd, Edinburgh, 558 pp.
- Favre A, Matuszak S, Sun H, Liu E, Yuan YM, Muellner-Riehl AN (2014) Two new genera of Gentianinae (Gentianaceae): *Sinogentiana* and *Kuepferia* supported by molecular phylogenetic evidence. Taxon 63(2): 342–354. <https://doi.org/10.12705/632.5>
- Favre A, Michalak I, Chen CH, Wang JC, Pringle JS, Matuszak S, Sun H, Yuan YM, Struwe L, Muellner-Riehl AN (2016) Out-of-Tibet: The spatio-temporal evolution of *Gentiana* (Gentianaceae). Journal of Biogeography 43(10): 1967–1978. <https://doi.org/10.1111/jbi.12840>
- Favre A, Pringle JS, Heckenauer J, Kozuharova E, Gao QB, Lemmon EM, Lemmon AR, Sun H, Tkach N, Gebauer S, Sun SS, Fu PC (2020) Phylogenetic relationships and sectional delineation within *Gentiana* (Gentianaceae). Taxon 69(6): 1221–1238. <https://doi.org/10.1002/tax.12405>
- Favre A, Pringle JS, Fu PC (2022) Phylogenetics support the description of a new sichuanese species, susanne's *Gentiana*, *Gentiana susanneae* (Gentianaceae). Systematic Botany 47(2): 506–513. <https://doi.org/10.1600/036364422X16512564801579>
- Fu PC, Twyford AD, Sun SS, Wang HY, Xia MZ, Tan CX, Zhou XJ, Chen SL (2021) Recurrent hybridization underlies the evolution of novelty in *Gentiana* (Gentianaceae) in the Qinhai-Tibetan Plateau. AoB Plants 13(1): plaa068. <https://doi.org/10.1093/aobpla/plaa068>
- Fu PC, Chen SL, Sun SS, Favre A (2022) Strong plastid degradation is consistent within section *Chondrophyllae*, the most speciose lineage of *Gentiana*. Ecology and Evolution 12(8): e9205. <https://doi.org/10.1002/ece3.9205>
- He D, Xiang X, He JS, Wang C, Cao G, Adams J, Chu H (2016) Composition of the soil fungal community is more sensitive to phosphorus than nitrogen addition in the alpine meadow on the Qinghai-Tibetan Plateau. Biology and Fertility of Soils 52(8): 1059–1072. <https://doi.org/10.1007/s00374-016-1142-4>
- Ho TN, Liu SW (1990) The infrageneric classification of *Gentiana* (Gentianaceae). Bulletin of the British Museum of Natural History 20: 169–192.
- Ho TN, Liu SW (2001) A worldwide monograph of *Gentiana*. Science Press, 455 pp.
- Ho TN, Liu SW (2010) Two new species of Gentianaceae from northwestern Yunnan, China. Novon: A Journal for Botanical Nomenclature 20(2): 166–169. <https://doi.org/10.3417/2003046>
- Ho TN, Pringle JS (1995) Gentianaceae. In: Wu ZY, Reven PH (Eds) Flora of China, Vol.16. Science Press, Beijing, China and Missouri Botanical Garden Press, St. Louis, Missouri, 140 pp.
- Ho TN, Liu SW, Wu QR (1988) Gentianaceae. In: Ho TN (Ed.) Flora Republicae Popularis Sinicae, vol.62. Science Press, Beijing, 411 pp.
- Ho TN, Liu SW, Lu XF (1996) A phylogenetic analysis of *Gentiana* (Gentianaceae). Journal of Systematics and Evolution 34(5): 505–530.
- Hsieh TY, Hsu TC, Kono Y, Ku SM, Peng CI (2007) *Gentiana bambuseti* (Gentianaceae), a new species from Taiwan. Botanical Studies 48(3): 349–355.
- Jiang M, Cui BW, Wu YL, Nan JX, Lian LH (2021) Genus *Gentiana*: A review on phytochemistry, pharmacology and molecular mechanism. Journal of Ethnopharmacology 264: 113391. <https://doi.org/10.1016/j.jep.2020.113391>

- Kalyaanamoorthy S, Minh B, Wong T, von Haeseler A, Jermiin L (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Mariana V, Ioan V, Andra VI (2013) Introducing in the architectural landscape of some *Gentiana* spontaneous species with ornamental. *Analele Universitatii din Oradea. Fascicula Protectia Mediului* 21: 308–315.
- Matuszak S, Favre A, Schnitzler J, Muellner-Riehl AN (2016) Key innovations and climatic niche divergence as drivers of diversification in subtropical Gentianinae in southeastern and eastern Asia. *American Journal of Botany* 103(5): 899–911. <https://doi.org/10.3732/ajb.1500352>
- Mayfield E (2021) Illustrated Plant Glossary. CSIRO Publishing, 321 pp. <https://doi.org/10.1071/9781486303540>
- Mirzaee F, Hosseini A, Jouybari BH, Davoodi A, Azadbakht M (2017) Medicinal, biological and phytochemical properties of *Gentiana* species. *Journal of Traditional and Complementary Medicine* 7(4): 400–408. <https://doi.org/10.1016/j.jtcme.2016.12.013>
- Ronquist F, Teslenko M, Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Shabir M, Kumar A, Dwivedi MD, Tiwari JK (2022) *Gentiana ranae* (Gentianaceae), a new species from India revealed by morphology and molecular phylogenetic analysis. *Rheedea* 32: 1–23. <https://doi.org/10.22244/rheedea.2021.32.01.01>
- Sun SS, Fu PC (2019) Study on taxonomy and evolution of Gentianeae (Gentianaceae). *Acta Botanica Boreali Occidentalis Sinica* 39(2): 363–370.
- Sun SS, Fu PC, Zhou XJ, Cheng YW, Zhang FQ, Chen SL, Gao QB (2018) The complete plastome sequences of seven species in *Gentiana* sect. *Kudoa* (Gentianaceae): Insights into plastid gene loss and molecular evolution. *Frontiers in Plant Science* 9: 493. <https://doi.org/10.3389/fpls.2018.00493>
- Wu LH, Leon CJ, Wang ZT (2012) *Gentiana zhenxiongensis* (Gentianaceae), a new species from Yunnan, China. *Annales Botanici Fennici* 49(3): 197–200. <https://doi.org/10.5735/085.049.0307>
- Yang SY, Du F, Wang J (2008) A new species of genus *Gentiana-Gentiana gracilis*. *Journal of Southwest Forestry College* 05: 1–2.
- Yang SY, Cao HF, Du F, Wang J, Zhang QR, Wang WT (2020) A new species of *Gentiana*, *Gentiana ciliolata* and its two subspecies of subsp. *ciliolata* and subsp. *longiloba* of Yunnan & Sichuan Provinces. *American Journal of Plant Sciences* 11(07): 1137–1143. <https://doi.org/10.4236/ajps.2020.117080>
- Yu JH, He H, Zhang BY, Yuan SS, Chen CM (2012) A new species of *Gentiana L.—G. hsin-ganica* from Inner Mongolica, China. *Bulletin of Botanical Research* 32(1): 1–3.
- Yuan YM, Küpfer P (1995) Molecular phylogenetics of the subtribe. Gentianinae (Gentianaceae) inferred from the sequences of internal transcribed spacers (ITS) of nuclear ribosomal DNA. *Plant Systematics and Evolution* 196: 207–226. <https://doi.org/10.1007/BF00982961>
- Yuan YM, Küpfer P (1997) The monophyly and rapid evolution of *Gentiana* sect. *Chondrophyllae* Bunge (Gentianaceae): Evidence from the nucleotide sequences of

the internal transcribed spacers of nuclear ribosomal DNA. Botanical Journal of the Linnean Society 123(1): 25–43. <https://doi.org/10.1111/j.1095-8339.1997.tb01403.x>

Yuan YM, Küpfer P, Doyle JJ (1996) Infrageneric phylogeny of the genus *Gentiana* (Gentianaceae) inferred from nucleotide sequences of the internal transcribed spacers (ITS) of nuclear ribosomal DNA. American Journal of Botany 83(5): 641–652. <https://doi.org/10.1002/j.1537-2197.1996.tb12750.x>

Appendix 1

Table A1. Species sequence information downloaded from the GenBank.

GenBank	Species	Voucher information	Herbarium
KU512339	<i>G. panthaica</i> Prain et Burk.	GXJ2011-055	
KT907686	<i>G. panthaica</i> Prain et Burk.	Favre & Matuszak 061a	KUN
KT907673	<i>G. mairei</i> Lev.	Wang hong et al. 01-0062	KUN
KT907643	<i>G. epichysantha</i> Hand.-Mazz.	Favre & Matuszak 131a	KUN
KF563953	<i>G. epichysantha</i> Hand.-Mazz.	Favre & Matuszak 131a	KUN
KT907728	<i>G. tatsienensis</i> Franch.	Wang Lisong, et al. 07-13	KUN
KU512348	<i>G. pseudoaquatica</i> Kusnezow	GXJ2011-043	
KT907644	<i>G. faucipilosa</i> H. Smith	Favre & Matuszak 37a	KUN
KF563954	<i>G. faucipilosa</i> H. Smith	Favre and Matuszak 37a	KUN
MT483837	<i>G. haynaldii</i> Kanitz		
MT483859	<i>G. haynaldii</i> Kanitz		
KT907613	<i>G. asterocalyx</i> Diels	Favre & Matuszak 106a	KUN
KT907653	<i>G. heleonastes</i> H. Smith ex Marq.	Favre 206a	KUN
KT907646	<i>G. forrestii</i> Marq.	Penghua, Liu Ende et al. 9538	KUN
KT907705	<i>G. rubicunda</i> Franch.	Wang hong et al. 03-1098	KUN
KU512351	<i>G. rubicunda</i> Franch.	YG2011392	
KT907651	<i>G. grata</i> H. Smith	Favre & Matuszak 32a	KUN
KT907666	<i>G. linoides</i> Franch. ex Hemsl.	Chen HS, CHC 2866	TNM
KU512331	<i>G. linoides</i> Franch. ex Hemsl.	GXJ2011-068b	
KT907687	<i>G. papillosa</i> Franck.	Chen HS, CHC 2867	TNM
KT907645	<i>G. flavomaculata</i> Hayata	Chen HS, CHC 2364	TNM
KT907618	<i>G. bella</i> Franch. ex Hemsl.	H.S. Chen CHC 2850	TNM
KT907741	<i>G. zollingeri</i> Fawcett	Chen HS CHC 2372	TNM
KU512311	<i>G. algida</i> Pall.	GXJ20130174	
KU512312	<i>G. algida</i> Pall.	PG110833	
KT907652	<i>G. handeliana</i> H. Smith	Liu Ende et al. 1209080	LZ
KT907680	<i>G. nubigena</i> Edgew.	Wang Lisong, et al. 07-55	KUN
KT907723	<i>G. striolata</i> T. N. Ho	Favre 221a	KUN
KU512317	<i>G. atunsiensis</i> W. W. Smith	GXJ2011-072	
KU512318	<i>G. atunsiensis</i> W. W. Smith	GXJ2011-080	
KT907627	<i>G. cephalantha</i> Franch. ex Hemsl.	Favre 325	KUN
KF563950	<i>G. cephalantha</i> Franch. ex Hemsl.	Favre 325	KUN
KT907642	<i>G. duclouxii</i> Franch.	Favre & Matuszak 076	KUN
MN124367	<i>G. lawrencei</i> var. <i>farreri</i> (I. B. Balfour) T. N. Ho	fu2016060_2	

GenBank	Species	Voucher information	Herbarium
MN124366	<i>G. lawrencei</i> var. <i>farreri</i> (I. B. Balfour) T. N. Ho	fu2016176_5	
KT907624	<i>G. caelestis</i> (Marq.) H. Smith	Favre & Matuszak 193a	KUN
KT907715	<i>G. sino-ornata</i> Balf.f.	Favre & Matuszak 224a	KUN
KU512319	<i>G. caelestis</i> (Marq.) H. Smith	GXJ2011-078	
KF563963	<i>G. purdomii</i> Marq.	Favre 311	KUN
KF563961	<i>G. phyllocayx</i> C. B. Clarke	Favre & Matuszak 33a	KUN
KT907690	<i>G. phyllocayx</i> C. B. Clarke	Favre & Matuszak 33a	KUN
MN339943	<i>G. stipitata</i> Edgew.		
MN339944	<i>G. stipitata</i> Edgew.		
AJ489922	<i>Swertia rosulata</i> (Baker) Klack.	Piso, Wohlhauser, Zeltner M023	NEU
MT199413	<i>Halenia taruga-gasso</i> Gilg	M005693166	
MK416129	<i>Gentianella gentianoides</i> (Franch.) H. Smith	xuechy090094	