

A new species of *Meryta* (Araliaceae) from the Marquesas Archipelago, French Polynesia

Frédéric Tronchet¹, Porter P. Lowry II^{1,2}

1 Département Systématique et Évolution (UMR 7205), Muséum National d'Histoire Naturelle, CP 39, 57 rue Cuvier, 75213 Paris CEDEX 05, France **2** Missouri Botanical Garden, P.O. Box 299, St. Louis, MO, 63166-0299, U.S.A.

Corresponding author: Porter P. Lowry II (Pete.Lowry@mobot.org)

Academic editor: Sandra Knapp | Received 18 April 2011 | Accepted 11 July 2011 | Published 12 July 2011

Citation: Tronchet F, Lowry II PP (2011) A new species of *Meryta* (Araliaceae) from the Marquesas Archipelago, French Polynesia. In: Lorence DH, Wagner WL (Eds) Botany of the Marquesas Islands: new taxa, combinations, and revisions. PhytoKeys 4: 149–156. doi: 10.3897/phytokeys.4.1408

Abstract

Meryta pastoralis F. Tronchet & Lowry, a new species from the island of Hiva Oa in the Marquesas archipelago, is described and illustrated. It differs from other Polynesian members of the genus by its fully free ovaries, a feature shared with one other species found in the region, *M. choristantha* (native to the Austral Islands), from which it can be distinguished by its ovate to spatulate (vs. elliptic to obovate) leaf shape. A preliminary risk of extinction assessment indicates that *M. pastoralis* is Critically Endangered.

Résumé

Une nouvelle espèce, *Meryta pastoralis* F. Tronchet & Lowry, de l'île d'Hiva Oa dans l'archipel des Marquises, est ici décrite et illustrée. Elle diffère des autres espèces du genre en Polynésie par ses ovaires totalement libres. Ce caractère est partagé uniquement par une autre espèce de la région – *M. choristantha* (endémique de l'archipel des Australes) – dont elle se distingue par ses feuilles à limbe ovale à spatulé (vs. elliptique à obovale). Une première estimation du statut de de risque d'extinction indique que l'espèce *M. pastoralis* est en danger critique.

Keywords

Meryta, Araliaceae, French Polynesia, Marquesas, Hiva Oa, conservation

Introduction

The genus *Meryta* J.R. Forst. & G. Forst. comprises a total of 37 species, 10 of which remain to be described (Frodin and Govaerts 2003, Tronchet and al. 2005a, Tronchet and Lowry unpubl. data), including the one proposed here. This distinctive group of monocaulous to well branched trees and shrubs is unique within Araliaceae in exhibiting a combination of simple leaves and a dioecious sexual system (Lowry 1988, Tronchet and al. 2005a). *Meryta* occurs primarily in the South Pacific, extending from New Zealand and New Caledonia in the west across to Henderson Island in the east, with a single member present north of the equator in the Caroline and Mariana Islands (Lowry 1993, Tronchet and al. 2005a). Without exception, each species is endemic to a single island or archipelago, and most are known from just one or a few populations. While they rarely comprise a significant component of the vegetation, the distinctive morphology of *Meryta* species makes them easy to recognize. A few taxa are cultivated, most notably *M. pauciflora* Hemsl. ex Cheesman, which is often listed in horticulture catalogues, and *M. sinclairii* (Hook. f.) Seem., widely grown in New Zealand and occasionally elsewhere.

A recent phylogenetic study based on molecular sequence data (Tronchet et al. 2005a) showed that *Meryta* is monophyletic, a finding that is consistent both with morphology and wood anatomy (Oskolski et al. 2007). Within Araliaceae, *Meryta* belongs to the *Polyscias-Pseudopanax* clade, as defined by Plunkett et al. (2004, 2005; see also Lowry et al. 2004), and within this clade it is most closely related to the SW Pacific genus *Plerandra*, which was recently expanded to include species from Melanesia long included in *Schefflera* (Lowry et al. in press). The molecular phylogeny of Tronchet et al. (2005a) also revealed two principal subclades within the genus, one comprising a majority of the species (including the type) and the other with two species, one each from New Zealand and Fiji.

As part of our taxonomic revision of *Meryta*, we describe here a new species endemic to the island of Hiva Oa in the Marquesas archipelago so that it can be included in the forthcoming *Vascular Flora of the Marquesas Islands* being prepared by D. H. Lorence and W. L. Wagner. This distinctive new entity, first collected by P.A. Schäfer in 1975, represents a significant range extension for the genus and a noteworthy addition to the flora of the Marquesas.

Systematics

Meryta pastoralis F. Tronchet & Lowry, sp. nov.

urn:lsid:ipni.org:names:77112742-1

http://species-id.net/wiki/Meryta_pastoralis

Figs 1-2

Haec species quoad inflorescentiae structuram ac ovaria omnino libera Merytae choristanthae Harms (ex Insulis Australibus) simillima, sed ab ea foliis anguste obovatis usque spathulatis (vs. late ellipticis usque subobovatis), carpellis 5 ad 11 (vs. 5 vel 6 tantum) atque fructu maturitate 7–9 × 7–10 (vs. 8–12 × 10–15) mm distinguitur; etiam quoad foliorum aspectum M. raiateensi J.W. Moore et aliquantum M. lanceolatae J.R. Forst. & G. Forst. similis, sed ab eis (atque adeo ab omnibus congeneris Societatis Insularum ac Tuamotu) ovariis omnino liberis distinguitur.

Type. French Polynesia. Marquesas Islands, Hiva-Oa, Hanamenu valley off Hanamenu trail, in *Metrosideros-Weinmannia-Dicranopteris linearis* wet forest, 09°47'50"S, 139°05'35"W, 908 m, 2 August 2005 (fr), S. Perlman 19767 (holotype: P [P00398408]!; isotypes, PAP!, PTBG!, US!).

Description. Monocaulous to sparsely branched tree, dioecious, Chamberlain architecture (tending toward Leeuwenberg architecture (Hallé 2004) when very old, fide P.A. Schäfer, 1975), 3–6(–10) m tall, without milky sap. Leaves simple, alternate, grouped at branch ends; petiole 1.5–2.5(–4) cm long, 2–3 mm in diam., base enlarged, slightly clasping, lenticels rarely present abaxially, without dark green transverse striations adaxially when fresh; ligule present adaxially at the base of the petiole, persistent, free portion triangular to widely triangular, 5–11(–17) mm long, margins entire, membranous, apex obtuse to acute; blade green, shiny adaxially when fresh, slightly lighter green abaxially, narrowly obovate to spatulate, 20.4–40 × 5.2–9.8 cm (l/w ratio 2.76–5.52), base symmetric to asymmetric, offset 2–5(–10) mm, mostly positively centered in adaxial view, attenuate, margin entire, undulate distally, minutely revolute, apex acute to obtuse, the extreme tip obtuse to mucronate, chartaceous to coriaceous, glabrous on both surfaces; venation brochidodromous, light green-yellow when fresh, with prominent arches lacking intramarginal veins; midvein strong and massive, straight, without evident abaxial thickenings; secondary veins (12–)17–22 pairs, diverging from the midvein by (53–)57–69° (in the widest part of the blade), the distal ones less divergent, weakly recurved near the midvein then strongly recurved toward the margin, joining with the next arch at an obtuse angle without forming a clear intra-marginal vein, intersecondary veins present; tertiary veins evident, not reaching the margin, straight to anastomosing at various angles, sometimes forming convex arcs in the intercostals zone; higher order veins visible in dry material, forming a fine, dense reticulum; veinlets forming quadri- to multi-angular areoles 0.5–1.5 mm in size. Juvenile foliage similar to adult, blade with an obtuse apex. Male material unknown (only old inflorescences seen). Female inflorescence terminal, erect, a raceme of spikes, with 1 degree of branching, axes and peduncle light green, bracts brown-red,




Figure 1. *Meryta pastoralis* F. Tronchet & Lowry. **A** branch with leaves and young infructescence **B** female flowers in bud subtended by bracts **C** female flowers at receptivity **D** young fruits. Line drawing by Alain Jouy from herbarium specimens; voucher Perlman 19767 (**A**, **D**), Schäfer 5922 (**B**), Perlman 10206 (**C**).

primary axis 15–20 cm long, 5–8 mm in diam. at the base when fresh, subtended by persistent or sometimes caducous triangular cataphylls 13–24 mm long, without lenticels, margins entire, mostly to almost entirely membranous, apex obtuse, bear-



Figure 2. *Meryta pastoralis* F. Tronchet & Lowry. Photo of female plant in fruit (Photo by Steve Perlman of Perlman 19767).

ing a dorsal, \pm foliaceous apicule; secondary axes 8–13(–17), 6–12 cm long (shorter distally), straight or sometimes slightly curved upward in the distal portion, diverging $35\text{--}60^\circ$ from the primary axis, each subtended by a caducous, triangular to narrowly triangular bract 7–20 mm long, without lenticels, margin denticulate, mostly membranous, apex acute to acuminate, sometimes apiculate; spikelets with 7–15(–27) flowers, the proximal one inserted 3–15 mm from the base of the secondary axis. Female flowers white when fresh, sessile; bractlets caducous or sometimes persistent, broadly triangular, 3–5 mm long, margin weakly dentate, membranous, brown-red, partially covered by the flower and later the fruit, apex acute to obtuse; petals 5–11, caducous in fruit, ovate, 1–2.5 mm long, weakly cucullate, recurved when flowers receptive, with a pronounced adaxial groove, apex acute; androecium present, isostemonous, filaments and anthers developed but pollen sacs empty, stamens with filaments 0.5–1 mm long, anthers 0.5 mm long; ovary inferior, (5–)7 or 8(–11)-carpellate, nectar disc epigynous, 2 mm diam., styles weakly differentiated, to 1 mm long, stigmas 1.5–3 mm long at receptivity, often strongly recurved to twisted. Fruit yellow-green when young, light purple at maturity, remains of petals and stamens brown, ovaries entirely free and distinct from one another; drupes globose to sub-globose, $7\text{--}9 \times 7\text{--}10$ mm, smooth and fleshy when fresh, ribbed when dry, the ribs corresponding to the 5–11 pyrenes.




Figure 3. Distribution map of *Meryta pastoralis* F. Tronchet & Lowry on Hiva Oa Island in the Marquesas archipelago (blue lines indicate primary water courses).

Distribution. Known only from the type locality in the Mt. Temetiu/Feani area at ca. 900 m elevation along the Hanamenu trail on the island of Hiva Oa (Fig. 3), the largest and highest in the Marquesas archipelago.

Ecology. *Meryta pastoralis* occurs on slopes and along streams in low stature, humid montane *Metrosideros-Weinmannia* forest at and above 900 m elevation.

Etymology. This species is named in honor of P.A. Schäfer, who was the first to collect it in 1975 while conducting botanical inventory work in the Marquesas, and who has contributed much to our knowledge of the flora of the archipelago. The name Schäfer means shepherd in Alsatian, which when translated into Latin provides the basis for our choice of the epithet *pastoralis*.

Conservation status. *Meryta pastoralis* is known from a single population along a wind-swept ridge below the highest point on Hiva Oa. With an Area of Occupancy of <10 km² and an Extent of Occurrence that is probably no larger, it meets the area requirements for Critically Endangered status under criteria B1 and B2 of the IUCN Red List Criteria (IUCN 2001). While the vegetation in the immediate area is still largely intact, increasing impacts from feral pigs, human disturbance and fire during the dry season (D. Lorence, pers. comm.) represent growing threats to *M. pastoralis*, prompting us to assign it a provisional threat status of Critically Endangered (CR B1ab(i,ii,iii,v) + 2ab(i,ii,iii,v)).

Other specimens examined. French Polynesia. Marquesas Islands, Hiva Oa, trail toward Hanamenu; in gulch off trail, low forest, 3 August 1988 (fr). S. Perlman 10206 (AD!, BISH! [2 sheets], F!, K!, MO!, NY!, P [P00372510]!, PAP!, PTBG!, US!); Teakatu, valley on north side of Hanamenu trail heading down to Hanamenu past summit crest, between Teakatau and Tepuna, along stream bottom in *Metrosideros-Weinmannia-Crossostylis* montane wet forest, [09°47'29"S, 139°05'41"W], 933 m, 26

August 1995 (st), S. Perlman & J.-Y. Meyer 14894 (BISH!, MO!, PAP!, PTBG!, US!); Feani area, Tepuna, in gulch to north of Hanamenu trail, down west side of summit crest to 1075 meters elev. and drop into gulch, in *Metrosideros-Weinmannia* forest, 09°47'49"S, 139°05'12"W, 900 m, 29 January 2003 (old ♂ fl), S. Perlman 18335 (BISH, P [P00398011]!, PAP, PTBG, US); Feani area, Tepuna, off Hanamenu trail on west side of summit crest, at 1075 meters elev. into gulch north side of trail, scattered plants in *Metrosideros-Weinmannia* forest, 09°47'49"S, 139°05'12"W, 29 January 2003 (♀ fl), S. Perlman 18336 (PAP, PTBG, US); Montagnes NW du Temetiu, entre la haute vallée de Hanamenu et la crête du Temetiu Feani, haute vallée, forêt humide, [09°48'00"S, 139°05'30"W], 925 m, 23 October 1975 (♀ fl), P.A. Schäfer 5922 (BISH!, K!, MO! [2 sheets], NTBGL!, P [P00372508 & P00372509]!, US!).

Discussion. Material of our new species most closely resembles specimens of *Meryta choristantha* Harms from the Austral Islands, with which it shares a similarly structured inflorescence and totally free ovaries, but differs in leaf shape (narrowly obovate to spatulate in *M. pastoralis* vs. widely elliptic to slightly obovate in *M. choristantha*), number of carpels (5–11 vs. 5 or 6), and fruit size at maturity (7–9 × 7–10 vs. 8–12 × 10–15 mm). The leaves of *Meryta pastoralis* are similar in appearance to those of *M. raiateensis* J.W. Moore, and to a lesser degree *M. lanceolata* J.R. Forst. & G. Forst., both from the Society Islands, but our new taxon differs from these entities, and indeed all other species in the Society Islands and on Tuamotu, by its totally free ovaries.

Results from ongoing molecular phylogenetic work (Tronchet et al. 2005a, 2005b, unpubl. data) suggest that *Meryta pastoralis* is part of a small clade now understood to comprise four species, also including *M. sinclairii* from New Zealand, *M. tenuifolia* A.C. Sm. from Fiji, and *M. choristantha*, which together correspond to *Meryta* sect. *Choristomeryta* Harms (Harms 1938). Material here assigned to *M. pastoralis* was referred to informally as '*M. schaeferi*' in Tronchet et al. (2005b).

Acknowledgments

We are grateful to David Lorence and Warren Wagner for suggesting that we publish this new species as part of the present series of novelties from the Marquesas Islands. We thank Alain Jouy for the fine line drawing, Steve Perlman for allowing us to use his color photo, Roy Gereau for translating the diagnosis into Latin, and David Lorence, Greg Plunkett and Warren Wagner for helpful comments on an earlier draft of the manuscript. This work was supported in part by grants from the U.S. National Science Foundation (DEB 0613728 to PPL) and the National Geographic Society (5793-96, to G.M. Plunkett).

References

- Frodin DG, Govaerts R (2003, publ. 2004) World Checklist and Bibliography of Araliaceae. The Royal Botanic Gardens, Kew, UK, 444 pp.

- Harms H (1938) Zur Kenntnis von *Meryta sonchifolia* Linden et André und einigen anderen Arten der Gattung. Notizblatt des botanischen Gartens und Museums zu Berlin-Dahlem 14: 315–321.
- Hallé (2004) Architecture des plantes. JPC Edition, Montpellier, 162 pages.
- IUCN (2001) IUCN Red List Categories and Criteria Version 3.1. Prepared by the IUCN Species Survival Commission. IUCN, Gland, Switzerland, 33 pp. [<http://www.iucnredlist.org/technical-documents/categories-and-criteria/2001-categories-criteria>]
- Lowry II PP (1988) A revision of Araliaceae from Vanuatu. Bulletin du Muséum National d'Histoire Naturelle, 4^e série, 11, 1989, sect. B, Adansonia 11: 117–155.
- Lowry II PP (1993) *Meryta* J.R. Forst. & G. Forst. Pp. 88–90 in van Balgooy MMJ (ed.), Pacific Plant Areas. Vol. 5. Rijksherbarium/Hortus Botanicus, Leiden University, The Netherlands.
- Lowry II PP, Plunkett GM, Wen J (2004) Generic relationships in Araliaceae: looking into the crystal ball. South African Journal of Botany 70: 382–392.
- Oskolski A, Kotina EL, Fomichev IV, Tronchet F, Lowry II PP (2007) Systematic implications of wood and bark anatomy in the Pacific island genus *Meryta* (Araliaceae). Botanical Journal of the Linnean Society 153: 363–379. [[http://onlinelibrary.wiley.com/doi/10.1111/j.1095-8339.2007.00612.x](http://onlinelibrary.wiley.com/doi/10.1111/j.1095-8339.2007.00612.x/abstract;jsessionid=6BD6EFAC497A22CFB3DC7038D811105C.d01t03)] doi:10.1111/j.1095-8339.2007.00612.x
- Plunkett GM, Wen J, Lowry II PP (2004) Intrafamilial classifications and characters in Araliaceae: Insights from the phylogenetic analysis of nuclear (ITS) and plastid (trnL-trnF) sequence data. Plant Systematics and Evolution 245: 1–39. [<http://www.ingentaconnect.com/content/schweiz/pde/2010/00000128/F0020001/art00004>] doi:10.1007/s00606-003-0101-3
- Plunkett GM, Lowry II PP, Frodin DG, Wen J (2005) Phylogeny and geography of *Schefflera*: pervasive polyphyly in the largest genus of Araliaceae. Annals of the Missouri Botanical Garden 92: 202–224. [<http://www.biodiversitylibrary.org/item/87371#page/208/mode/1up>]
- Tronchet F, Plunkett GM, Jérémie J, Lowry II PP (2005a) Monophyly and major clades of *Meryta* (Araliaceae). Systematic Botany 30: 657–670. [<http://www.ingentaconnect.com/content/aspt/sb/2005/00000030/00000003/art00016>] doi:10.1600/0363644054782279
- Tronchet F, Plunkett GM, Jérémie J, Lowry II PP (2005b) Phylogeny and biogeography of the Pacific genus *Meryta* (Araliaceae) based on ITS and ETS sequence data. Poster presented at the 17th International Botanical Congress, Vienna, Austria.