

Resurrection of the genus *Staphisagria* J. Hill, sister to all the other Delphinieae (Ranunculaceae)

Florian Jabbour, Susanne S. Renner

Systematic Botany and Mycology, University of Munich (LMU), Menzinger-Str. 67, 80638 Munich, Germany

Corresponding author: Florian Jabbour (jabbour@bio.lmu.de)

Academic editor: Th. Couvreur | Received 5 September 2011 | Accepted 12 October 2011 | Published 29 November 2011

Citation: Jabbour F, Renner SS (2011) Resurrection of the genus *Staphisagria* J. Hill, sister to all the other Delphinieae (Ranunculaceae). *PhytoKeys* 7: 21–26. doi: 10.3897/phytokeys.7.2010

Abstract

Molecular sequence data show that the three species of *Delphinium* subg. *Staphisagria* (J. Hill) Peterm. form the sister clade to *Aconitum* L., *Aconitella* Spach, *Consolida* (DC.) S.F. Gray, and all remaining species of *Delphinium* L. To account for this finding we resurrect *Staphisagria* J. Hill (1756). Names in *Staphisagria* are available for two of the species. We here make the required new combination for the third species, *Staphisagria picta* (Willd.) F. Jabbour, provide a key to the species, and illustrate one of them.

Keywords

Aconitum, *Delphinium*, Mediterranean region, molecular phylogeny, nomenclature, *Staphisagria*

Introduction

Delphinium staphisagria L., *D. requienii* DC., and *D. pictum* Willd. are annual or biennial species of the tribe Delphinieae (Ranunculaceae) that occur in the Mediterranean basin (see ‘Nomenclature and key to the species’ section for a more detailed description of their distribution areas). They are considered endangered (Olivier et al. 1995, Fraga et al. 2004) because of changing land use patterns and bottlenecks caused by irregular demography (Orellana et al. 2009a). All three species are protected in France (Olivier et al. 1995).

Linnaeus knew only *D. staphisagria*, which he described as *Delphinium nectariis diphyl-lis, foliis palmatis lobis integris*. With the recognition in the early 19th century that there were two additional species resembling *D. staphisagria*, Spach (1839) grouped all three

in the genus *Staphisagria* established by John Hill in 1756 for Linnaeus's *D. staphisagria*. Spach's ranking of the three species as a separate genus, distinct from *Delphinium*, however, gained few followers and no modern treatment appears to have accepted *Staphisagria*.

Molecular phylogenetic studies of the Delphinieae recently revealed that the three species of *Staphisagria sensu* Spach are the sister clade to all other Delphinieae (Maximum Likelihood bootstrap support: 90%), a group of 650-700 species ranging from Eurasia into North America and with a few isolated species on West and East African mountains (Jabbour and Renner, unpublished data; Fig. 1). This discovery fits with several characters of the three species that are unusual in *Delphinium*. For example, *Staphisagria* species have eight chromosome pairs of staggered size (see Fig. 3 in Verlaque and Aboucaya 2001), while most *Delphinium* have a bimodal karyotype of two long and six short chromosome pairs (Gregory 1941; Kurita 1955; Blanché and Simon 1987; Yang 1996, 2001). Species of *Aconitum* subg. *Lycocotonum* (c. 50 species) and the three species of *Staphisagria* (Verlaque and Aboucaya 2001) have a similar karyotype, suggesting parallel chromosomal reconfigurations. The *Staphisagria* species also have C19 aconitine-type alkaloids (De La Fuente and Reina 1990) and *Aconitum*-like stomata and pollen (Blanché 1991). Flowers of *Staphisagria* are less zygomorphic than those of the remaining Delphinieae and their nectar spurs are only 2–7 mm long (Bosch 1997; Verlaque and Aboucaya 2001). This last trait probably reflects predominant self-fertilization, with reduced reliance on nectar-foraging bees for cross-pollination (Bosch 1997; Bosch et al. 2001).

To account for the phylogenetic relationships in the Delphinieae (Fig. 1), we here resurrect the genus *Staphisagria* and make the required new combination for a species for which Spach (1839) did not provide a legitimate name.

Nomenclature and key to the species

Staphisagria J. Hill, Brit. Herbal: 44. 1756.

Type. *Delphinium staphisagria* L., Sp. Pl.: 531. 1753 [original type, cited by its nomen specificum legitimum].

Key to the species

- | | | |
|---|--|-----------------------|
| 1 | Spur of the upper tepal 1/5–1/3 as long as perianth segments (Fig. 2C); seeds 5.5–7.5 mm (Fig. 2F)..... | <i>S. macrosperma</i> |
| – | Spur of the upper tepal at least 2/5 as long as perianth segments; seeds 3–4.5 mm..... | 2 |
| 2 | Inflorescence axis, pedicels, and outside of perianth segments shortly pubescent; bracteoles inserted at the base of the pedicels..... | <i>S. picta</i> |
| 2 | Inflorescence axis, pedicels, and outside of perianth segments villose-hirsute; bracteoles inserted at some distance above the base of the pedicels... <i>S. requienii</i> | |




Figure 1. Schematic representation of the phylogenetic relationships in the Delphinieae **A** as suggested by studies anterior to the findings of Jabbour and Renner (unpublished data) and **B** as found with combined nuclear (ITS) and plastid (*trnL* intron and adjacent *trnL-trnF* intergenic spacer) DNA sequences (2088 aligned nucleotides) from 191 accessions representing 185 species of Delphinieae (Jabbour and Renner, unpublished data). In this study, taxon sampling covered all genera and subgenera of the tribe. Numbers above branches indicate Maximum Likelihood bootstrap supports.

Staphisagria [“*Staphysagria*”] *macroisperma* Spach, Hist. Nat. Vég. 7: 348. 1839.

Replaced name. *Delphinium staphisagria* L., Sp. Pl.: 531. 1753. TYPE: LINN 694/12, Habitat in Istria, Dalmatia, Calabria, Apulia, Creta, Galloprovincia [South France]. The geographic origin of Linneaus's type cannot be narrowed down (Munz 1967, Iltarlsan 1996, Jarvis 2007).

Herbarium specimen studied: Greece: Crete, Nomos Lassithiou, ravine between Zákros and Kato Zákros, 70 m, 15 May 2002, E. Vitek 02-205 (W, GZU, M, MA).

Comments. Spach had to chose a new name for this Linnaean species because *S. staphisagria* would be an exact tautonym (not permitted in botanical nomenclature), and since his misspelling of Hill's genus (as *Staphysagria*) is a correctable error (variant spelling), Spach's name *S. macroisperma* is legitimate. Among the three species of the genus *Staphisagria*, *S. macroisperma* has the largest distribution. Because of its ancient use in medicine (Cristofolini and Mossetti 1998), it is found all around the Mediterranean basin (Greuter et al. 1989; Orellana et al. 2009a). It grows in rocky areas, and is adapted to nitrophilous and disturbed habitats (Orellana et al. 2009a). Figure 2 shows key morphological characteristics of *S. macroisperma*.




Figure 2. *Staphisagria macrosperma* **A** young plant with the cotyledons and two palmate leaves **B** young inflorescence with floral buds subtended by a bract and bracteoles **C** side view of a flower showing the very short spur (or bulge) on the dorsal petaloid tepal **D** three-quarter view of a flower showing four developed staminodes: two lateral (black arrows) and two spurred (white arrows). The tips of the spurs are nested within the bulge of the dorsal tepal **E** three follicles **F** Gravity-dispersed poisonous seeds (c. 6 mm in length). Scale bars: 1 cm.

***Staphisagria requienii* (DC.) Spach, Hist. Nat. Vég. 7: 350. 1839.**

Basionym. *Delphinium requienii* DC., Fl. Franç. (DC. & Lamarck), ed. 3. 5: 642. 1805.

Herbarium specimen studied: France: Var, Hyères, Porquerolles island, 12 Jun 1961, Gavelle s.n. (M).

Comments. *Staphisagria requienii* is a narrow endemic of the Mediterranean Archipelago of Hyères, Var, South of France (Verlaque et al. 1991). It grows in a variety of habitats, like crops, calcareous rocks, and degraded areas along roads (Orellana et al. 2009b).

***Staphisagria picta* (Willd.) F. Jabbour, comb. nov.**

Basionym. *Delphinium pictum* Willd., Enum. Pl. [Willdenow] 1: 574. 1809. SYNTYPES: Röpert, D. (Ed.) 2000- (continuously updated): Digital specimen images at the Herbarium Berolinense. Published on the Internet <http://ww2.bgbm.org/herbarium/> Barcode: B -W 10324 -01 0 / ImageId: 164585) and Barcode: B -W 10324 -02 0 / ImageId: 164601) [accessed 02-Sept-11].

Herbarium specimen studied: Balearic Islands: Majorca, Puntas de Covas, top of sea cliffs, amongst limestone boulders, 100 m, April 1988, F.J. Rumsey s.n. (M).

Comments. The new combination is necessary because *S. brevipes* Spach (1839) is illegitimate since it included the older name *D. pictum*. *Staphisagria picta* is endemic to Corsica, Majorca, and Sardinia. Its main habitats are open grasslands covering rocky places from 150 up to 600 m above sea level (Orellana et al. 2009b)

Acknowledgements

We thank Werner Greuter for nomenclatural advice and the anonymous reviewer who made important comments and suggestions. Financial support for FJ's project came from the German Science Foundation (DFG) grant no. RE 603/12-1.

References

- Blanché C (1991) Revisió biosistemàtica del gènere *Delphinium* L. a la Peninsula Ibèrica i a les illes Balears. Arxius de la Secció de Ciències, Institut d'Estudis Catalans, Barcelona (1991): 1–288.
- Blanché C, Simon J (1987) Données cytotaxonomiques sur les *Consolida* (DC.) S. F. Gray (Ranunculaceae) en Méditerranée occidentale. Saussurea 18: 1–10.
- Bosch M (1997) Biologia de la reproducció de la tribu Delphinieae a la Mediterrània Occidental. PhD thesis. Institut d'Estudis Catalans. Arxius de les Seccions de Ciències, CXX, Secció de Ciències Biològiques.
- Bosch M, Simon J, Molero J, Blanché C (2001) Breeding systems in tribe Delphinieae (Ranunculaceae) in the western Mediterranean area. Flora 196: 101–113.

- Cristofolini G, Mossetti U (1998) Interpretation of plant names in a late medieval medical treatise. *Taxon* 47: 305–319. doi: 10.2307/1223760
- De La Fuente G, Reina M (1990) Some phytochemical studies of the genera *Aconitum* L., *Delphinium* L. and *Consolida* (DC.) Gray. *Collectanea Botanica* (Barcelona) 19: 129–140.
- Fraga P, Mascaró C, Carreras D, Garcia O, Pallicer X, Pons M, Seoane M, Truyol M (2004) Catàleg de la flora vascular de Menorca. Collecció Recerca N°9. Institut Menorquí d'Estudis, Menorca.
- Gregory W (1941) Phylogenetic and cytological studies in the Ranunculaceae. *Transactions of the American Philosophical Society, Series 2* 31: 443–521.
- Greuter W, Burdet H, Long G (1989) Med-Checklist, vol. 4 (Lauraceae-Rhamnaceae). Editions des Conservatoire et Jardin Botaniques de Genève, Geneva.
- Hill J (1756) Genus XI. Stavesacre. *Staphisagria*. In: Osborne T, Shipton J, Hodges J (Eds) *The British herbal: an history of plants and trees, natives of britain cultivated for use, or raised for beauty*, 44.
- Ilarslan R (1996) A revision of the genus *Delphinium* L. (Ranunculaceae) in Turkey. *Turkish Journal of Botany* 20: 133–161.
- Jarvis C (2007) *Delphinium staphisagria*. In: Order out of chaos: Linnean plant names and their types. Linnean Society of London, in association with the Natural History Museum, London, 478.
- Kurita M (1955) Cytological studies in Ranunculaceae, III. The karyotypes of several species in *Delphinium*, *Lycocotonum* and *Aconitum*. *Botanical Magazine Tokyo* 68: 248–251.
- Munz P (1967) A synopsis of the Asian species of *Delphinium*, sensu stricto. *Journal of the Arnold Arboretum* 48: 249–302.
- Olivier L, Galland J, Maurin H, Roux J (1995) *Livre Rouge de la Flore Menacée de France*. Tome I: Espèces Prioritaires. Muséum National d'Histoire Naturelle. Institut d'Ecologie et de Gestion de la Biodiversité. Service du Patrimoine Naturel. Collection Patrimoines Naturels, vol. 20, Série Patrimoine Génétique, Paris, 4.
- Orellana M, López-Pujol J, Blanché C, Rovira A, Bosch M (2009a) Genetic diversity in *Delphinium staphisagria* (Ranunculaceae), a rare Mediterranean dysploid larkspur with medicinal uses. *Genetica* 135: 221–232. doi: 10.1007/s10709-008-9271-9
- Orellana M, Blanché C, Simon J, Bosch M (2009b) Genetic diversity within and among disjunct populations of the Mediterranean Island endemic *Delphinium pictum* and *D. regnieri* (Ranunculaceae). *Folia Geobotanica* 44: 47–63. doi: 10.1007/s12224-009-9028-y
- Spach E (1839) *Histoire naturelle des végétaux. Phanérogames*. Tome septième. 1–538.
- Verlaque R, Aboucaya A (2001) Position des *Delphinium* du sous-genre *Staphisagria* dans la tribu des Delphinieae Warm (Ranunculaceae). *Bocconeia* 13: 189–200.
- Verlaque R, Aboucaya A, Cardona A, Contandriopoulos J (1991) Quelques exemples de spéciation insulaire en Méditerranée occidentale. *Botanika Chronika* 10: 137–153.
- Yang Q (1996) A karyotype study of 15 species in the tribe Delphinieae (Ranunculaceae) from China. *Acta Phytotaxonomica Sinica* 34: 39–47.
- Yang Q (2001) Cytology of 12 species in *Aconitum* L. and of 18 species in *Delphinium* L. of the tribe Delphinieae (Ranunculaceae) from China. *Acta Phytotaxonomica Sinica* 39: 502–514.