

Taxonomic studies of *Glochidion* (Phyllanthaceae) from the Indo-China Peninsula (I): *G. shanense*, a new species from Myanmar

Gang Yao¹, Jie Cai², Youheng Wu^{3,4}, Xuefei Yang^{5,6}, Thaung Naing Oo⁷,
Aung Zaw Moe⁷, Shixiao Luo³

1 South China Limestone Plants Research Centre, College of Forestry and Landscape Architecture, South China Agricultural University, Guangzhou 510642, China **2** Plant Germplasm and Genomics Centre, Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, 650201, China **3** Key Laboratory of Plant Resources Conservation and Sustainable Utilisation, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 516650, China **4** University of Chinese Academy of Sciences, Beijing 1000049, China **5** Key Laboratory of Economic Plants and Biotechnology and Yunnan Key Laboratory for Wild Plant Resources, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China **6** Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences, Yezin, Nay Pyi Taw 05282, Myanmar **7** Forest Research Institute, Forest Department, Ministry of Environmental Conservation and Forestry, Yezin, Nay Pyi Taw 05282, Myanmar

Corresponding author: Shixiao Luo (luoshixiao@scbg.ac.cn)

Academic editor: A. Sennikov | Received 16 February 2018 | Accepted 13 March 2018 | Published 27 March 2018

Citation: Yao G, Cai J, Wu Y, Yang X, Oo TN, Moe AZ, Luo S (2018) Taxonomic studies of *Glochidion* (Phyllanthaceae) from the Indo-China Peninsula (I): *G. shanense*, a new species from Myanmar. *PhytoKeys* 96: 57–62. <https://doi.org/10.3897/phytokeys.96.24497>

Abstract

Based on morphological studies performed on live plants in the field and specimens deposited in herbaria, a new species, *Glochidion shanense* Gang Yao & Shixiao Luo (Phyllanthaceae, Phyllanthaceae), is here described and illustrated. The species is morphologically most similar to *G. ellipticum* Wight, but can be distinguished from the latter by having hairy branchlets, longer pedicels, uniseriate and narrowly triangular sepals of female flowers, 4–5-locular ovaries, stout and cylindric persistent style on fruits.

Keywords

Phyllanthaceae, *Glochidion*, Morphology, Myanmar

Introduction

Glochidion J.R. et G. Forst. is the second largest genus in the tribe Phyllanthae Dumortier, Phyllanthaceae Martynov (Webster 1994). It consists of more than 300 species mainly distributed in the Indo-Pacific, east to southeast Polynesia and south into Australia (Govaerts et al. 2000, Li and Gilbert 2008) and can be distinguished from other members in Phyllanthae by the absence of a floral disc, apiculate anthers, usually confluent styles and fleshy seed-coat (Li 1994, Webster 1994).

Recently, the leafflower plant and leafflower moth system, a new mutualism model system that is similar to the fig-fig wasp and yucca-yucca moth pollination systems, was described between Phyllanthae plants and *Epicephala* moth (Kato et al. 2003, Kawakita et al. 2015, Luo et al. 2017) and it provides a new opportunity to study the mechanisms of biodiversity development and maintenance. *Glochidion* is the largest host plant lineage within this newly described mutualism system, in which at least five host plant lineages were identified (Kawakita and Kato 2009). However, a comprehensive taxonomic study of *Glochidion* is still lacking, especially for relevant species distributed in the Indo-China Peninsula, since the latest comprehensive taxonomic investigation of *Glochidion* from there can be dated back to Beille (1927). Recently, taxonomic studies of *Glochidion* species from Thailand (van Welzen 2007) and Vietnam (Nguyen 2007) have been conducted. In Myanmar, Kurz (1877) accepted 14 *Glochidion* species in Forest Flora of British Burma and Kress et al. (2003) recorded 33 *Glochidion* species in A Checklist of the Trees, Shrubs, Herbs and Climbers of Myanmar.

During the fieldwork in Shan State, eastern Myanmar, in December 2015, two of the authors (J. Cai and X.F. Yang) collected some Phyllanthae specimens and one belongs to *Glochidion*, which superficially differs from congeneric taxa recorded in Myanmar and its adjacent countries. After a detailed morphological investigation and herbaria examination for all the *Glochidion* species recorded in this region, it was confirmed that the species is new to science, thus it is formally described below. The new species belongs to section *Glochidion*, which is characterised by having three stamens in male flowers (Li 1994).

Materials and methods

In addition to fieldwork in mountain areas of the eastern Myanmar, the present study also included analyses of *Glochidion* material from herbaria HITBC, IBSC, K, KUN, P, PE and US, as well as consideration of the taxonomic literature of China (Li 1994, Li and Gilbert 2008), India (Hooker 1887, Balakrishnan and Chakrabarty 2007, Chakrabarty and Balakrishnan 2009), Indo-China Peninsula (Beille 1927), Myanmar (Kurz 1877, Kress et al. 2003), Thailand (van Welzen 2007) and Vietnam (Nguyen 2007). Morphological analyses in the present study were performed on live plants in the field and specimens deposited in herbaria. Herbarium abbreviations follow the Index Herbarium (Thiers 2018+).

Taxonomy

Glochidion shanense Gang Yao & Shixiao Luo, sp. nov.

urn:lsid:ipni.org:names:77177610-1

Figures 1–2

Diagnosis. The species is similar to *G. ellipticum* Wight in morphology, but differs from the latter by its branchlets pubescent, pedicels of female flowers 1.5–2 mm long, sepals of female flowers uniseriate and narrowly triangular, apex of style truncate slightly, persistent style of fruit stout and cylindric.

Type. MYANMAR. Shan State, Pindaya, near the Htwet Ni village, West of Pindaya town, at an elevation of 1396 m, forest understory, in flowering and fruiting, 25 December 2015, *Jie Cai et al.* 15CS10794 (holotype, KUN!; isotype, KUN!, IBSC!)

Description. Shrubs or treelets, up to 2 m; monoecious; branchlets pubescent. Leaf blade oblong or elliptic, 9–13.5 × 4.5–6.5 cm, papery, slightly leathery, with apex acuminate to round and base broadly cuneate, sparsely pubescent along veins adaxially, pubescent abaxially; midvein and 6–9-paired lateral veins elevated abaxially. Petiole 3–4 mm long, pubescent. Stipules narrowly triangular, 2–4 mm long, pubescent. Male flowers: pedicels 6–10 mm long, densely tomentose; sepals 6, biseriate, oblong or ovate, densely tomentose; stamens 3. Female flowers: in axillary clusters, pedicels 1.5–2 mm long, densely strigose; sepals 6, uniseriate, narrowly triangular, densely strigose; ovary depressed globose, 4–5-locular, densely strigose; style connate into a cylinder, ca. 1 mm long, slightly truncate at apex, apex 4–5-lobed. Capsules depressed globose, 8–9 mm in diameter, ca. 4 mm high, sparsely pubescent, 8–10-grooved; persistent style cylindric, ca. 1 mm long; fruiting pedicels 4–5 mm long, stout, densely pubescent; seed laterally compressed, orange.

Distribution and habitat. This new species is currently known only from its type locality, Shan State, eastern Myanmar, where it grows in the broadleaved and coniferous dry forest dominated by *Docynia indica* (Wall.) Decne., *Schima wallichii* (DC.) Korth. and *Pinus* species.

Etymology. *Glochidion shanense* is named after its type locality, Shan State in eastern Myanmar.

Taxonomic discussion. The species resembles *Glochidion ellipticum* Wight, a species widely distributed from eastern Himalaya to Taiwan Island, but differs from the latter by its branchlets densely pubescent (Figure 2G), pedicels of male flowers tomentose, pedicels of female flowers densely strigose and up to 1.5–2 mm long (Figures 2A, D), sepals of female flower uniseriate and narrowly triangular in shape (Figures 2A, D), ovaries 4–5-locular, style cylindric and truncate at apex (Figure 2D), fruits 8–9 mm in diameter and 8–10-grooved (Figure 2G), persistent styles cylindric and ca. 1 mm long (Figure 2G). In contrast, *G. ellipticum* has the branchlets glabrous (Figures 2B, H), pedicels of male flowers glabrous (Figure 2H), female flowers sessile (Figures 2B, E), sepals of female flowers biseriate and oblong in shape (Figure 2B), ovaries 3–4 (5)-locular, style columnar to columnar-conical (Figures 2B, E), fruits 6–8 (rarely up to 10) mm in diameter and shallowly 6–8 (rarely 10)-grooved (Figure 2H), persistent styles obscure or slightly elevated but far less than 1 mm long (Figure 2H).




Figure 1. Line drawing of *Glochidion shanense* Gang Yao & Shixiao Luo, sp. nov. **A** Habit **B** Female flower **C** Male flower **D** Fruit **E** Seed. Draw by Ling Wang on *Jie Cai et al. 15CS10794* (KUN).




Figure 2. Morphological comparisons between *Glochidion ellipticum* Wight (**B, C, E, H**) and *G. shanense* Gang Yao & Shixiao Luo (**A, D, F, G**). **A, B, D, E** Female flower **C, F** Branchlets **G, H** Fruit.

Acknowledgments

The authors are grateful to the Forest Research Institute of Myanmar for permission and support in conducting this study in eastern Myanmar. We are indebted to Mrs. Ling Wang in Kunming Institute of Botany, Chinese Academy of Sciences, for preparing the illustration. This research was financially supported by a project of the Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences (Grant No. Y4ZK111B01).

References

- Balakrishnan NP, Chakrabarty T (2007) The family Euphorbiaceae in India: A synopsis of its profile, taxonomy and bibliography. Bishen Singh and Mahendra Pal Singh, Dehradun, 500 pp.
- Beille L (1927) *Glochidion*. In: Lecomte H (Ed.) Flore générale de l'Indo-Chine 5. Masson, Paris, 608–628.
- Chakrabarty T, Balakrishnan NP (2009) The family Euphorbiaceae in Sikkim State, India. *Journal of Economic and Taxonomic Botany* 33: 483–539.
- Govaerts R, Frodin DG, Radcliffe-Smith A, Carter S (2000) World checklist and bibliography of Euphorbiaceae (Pandaceae). Royal Botanic Garden, Kew, UK, 1622 pp.
- Hooker JD (1887) The Flora of British India 5. Macmillan, London, 910 pp.
- Li PT (1994) Flora Reipublicae Popularis Sinicae 44. Science Press, Beijing, 217 pp.
- Li PT, Gilbert MG (2008) *Glochidion*. In: Wu ZY, Raven PH (Eds) Flora of China 11. Science Press & Missouri Botanical Garden Press, Beijing & St. Louis, 193–202.
- Luo SX, Yao G, Wang ZW, Zhang DX, Hembry DH (2017) A novel, enigmatic basal leafhopper moth lineage pollinating a derived leafhopper host illustrates the dynamics of host shifts, partner replacement, and apparent coadaptation in intimate mutualisms. *The American Naturalist* 189: 422–435. <https://doi.org/10.1086/690623>
- Kato M, Takimura A, Kawakita A (2003) An obligate pollination mutualism and reciprocal diversification in the tree genus *Glochidion* (Euphorbiaceae). *Proceedings of the National Academy of Sciences of the United States of America* 100: 5264–5267. <https://doi.org/10.1073/pnas.0837153100>
- Kawakita A, Kato M (2009) Repeated independent evolution of obligate pollination mutualism in the Phyllentheae-*Epicephala* association. *Proceedings of the Royal Society of London B: Biological Sciences* 276: 417–426. <https://doi.org/10.1098/rspb.2008.1226>
- Kawakita A, Mochizuki K, Kato M (2015) Reversal of mutualism in a leafhopper–leafhopper moth association: the possible driving role of a third-party partner. *Biological Journal of the Linnean Society* 116: 507–518. <https://doi.org/10.1111/bij.12633>
- Kress J, Robert A, DeFilippes E, Kyi YY (2003) A Checklist of the Trees, Shrubs, Herbs, and Climbers of Myanmar. <http://www.botany.si.edu/myanmar> [Accessed 3 February 2018]
- Kurz S (1877) Forest Flora of British Burma 2. Office of the superintendent of government printing, Calcutta, 613 pp.
- Nguyen NT (2007) Taxonomy of Euphorbiaceae in Vietnam. Vietnam National University Publishers, Hanoi, 407 pp.
- Thiers B (2018+) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. The New York Botanical Garden, New York. Available from: <http://sweetgum.nybg.org/ih/> [Accessed: 3 February 2018]
- van Welzen PC (2007) *Glochidion*. In: Chayamarit K, van Welzen PC (Eds) Flora of Thailand 8(2). Royal Forest Department, Bangkok, 308–331.
- Webster GL (1994) Synopsis of the genera and suprageneric taxa of Euphorbiaceae. *Annals of the Missouri Botanical Garden* 81: 33–144. <http://www.jstor.org/stable/2399909>