

Taxonomic studies on the genus *Isotrema* (Aristolochiaceae) from China: II. *I. brevilimbium* (Aristolochiaceae), a new species from Guizhou, China

Jun Wang¹, Ji-Dong Ya³, Cheng Liu³, Guang Liu⁴, Feng Cao⁵, Jin-Shuang Ma²,
Xin-Xin Zhu¹

1 College of Life Sciences, Xinyang Normal University, Xinyang, 464000, Henan, China **2** Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, Shanghai Chenshan Botanical Garden, Shanghai 201602, Shanghai, China **3** Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, China **4** Haizhu District Experimental Primary School, Guangzhou 510245, Guangdong, China **5** Weining 553100, Guizhou, China

Corresponding author: Jin-Shuang Ma (jinshuangma@gmail.com); Xin-Xin Zhu (zhuzhu8niuniu@126.com)

Academic editor: Elton John de Lirio | Received 5 March 2020 | Accepted 1 June 2020 | Published 3 July 2020

Citation: Wang J, Ya J-D, Liu C, Liu G, Cao F, Ma J-S, Zhu X-X (2020) Taxonomic studies on the genus *Isotrema* (Aristolochiaceae) from China: II. *I. brevilimbium* (Aristolochiaceae), a new species from Guizhou, China. *PhytoKeys* 152: 15–25. <https://doi.org/10.3897/phytokeys.152.51760>

Abstract

A new species of *Isotrema* was recently discovered from Guizhou, China and is here named as *I. brevilimbium*. It is most similar to *I. ovatifolium* and *I. wardianum*, but differs in the morphology of leaves and flowers. A detailed description for the new species, along with line drawings, photographs, as well as morphological comparisons with similar species, are provided. In addition, the distribution of *I. wardianum* in China is here confirmed.

Keywords

Aristolochia wardiana, morphology, subgenus *Siphisia*, taxonomy

Introduction

Isotrema Raf. (Aristolochiaceae), previously treated as a subgenus of *Aristolochia* L., was recently reinstated as an independent genus based on molecular and morphological evidence (Zhu et al. 2019a). It can be distinguished from *Aristolochia* by the follow-

ing set of characters: perianth strongly curved, gynostemium 3-lobed, anthers paired on the outer surface of each gynostemium segment, and capsule dehiscing basipetally (Do et al. 2015a; Zhu et al. 2019a). Several new species of *Isotrema* have been found and described from China and its neighbouring countries during recent years (Liu and Deng 2009; Xu et al. 2011; Yao 2012; Huang et al. 2013, 2015; Wu et al. 2013, 2015; Do et al. 2014, 2015a, 2015b, 2015c, 2015d, 2016, 2017, 2019; Huang et al. 2014; Lu and Wang 2014; Ohi-Toma et al. 2014; Zhu et al. 2015, 2016, 2017a, 2017b, 2018, 2019b, 2019c; Gong et al. 2018; Yang et al. 2018; Li et al. 2019; Peng et al. 2019; Zhou et al. 2019; Cai et al. 2020a, 2020b). Currently, a total of 106 species have been reported from *Isotrema*, most of which are distributed in eastern and southern Asia, with some species further extended to northern and central America (Zhu et al. 2019a). China accommodates ca. 66 species, among which 55 species are endemic (Huang et al. 2003; Li et al. 2019; Peng et al. 2019; Zhou et al. 2019; Zhu et al. 2019a, 2019b, 2019c, 2019d; Cai et al. 2020a, 2020b).

During our recent field explorations to southern China, an unknown species of *Isotrema* was collected. Our subsequent examination of specimens from 39 public herbaria (A, BM, BR, CDBI, CSFI, CSH, E, EMA, GXMI, HAST, HENU, HHBG, HIB, HITBC, HNWP, IBK, IBSC, K, KYO, KUN, L, LBG, LE, NAS, NTUF, P, PE, PEM, SM, SNU, SYS, TAI, TI, W, WCU, WU, WUK, XYTC, YUKU; abbreviations follow Thiers 2020) and study of related literature (Hwang 1981, 1988; Ma 1989a, 1989b; Tao 1997; Huang et al. 2003; Do et al. 2015a; Do and Nghiem 2017; Yang et al. 2018; Zhu et al. 2019a, 2019d) suggested it to be a new species. Hereafter, we describe it as *I. brevilimbus* X.X.Zhu, Jun Wang & F.Cao. Moreover, *I. wardianum* (J.S. Ma) X.X. Zhu, S. Liao & J.S. Ma was recently published (Zhu et al. 2019a) based on its basionym *A. wardiana* J.S. Ma, previously only known from Myanmar and India (Ma 1989a), which was recently collected from Medog County, Tibet, and here its distribution in China is confirmed. Measurements and morphological characters of *I. brevilimbus*, *I. ovatifolium* and *I. wardianum* were made from both dried specimens and field observations of living plants, as well as including related literature. The morphological characters of the description follow Huang et al. (2003).

Taxonomy

***Isotrema brevilimbus* X.X.Zhu, Jun Wang & F.Cao, sp. nov.**

urn:lsid:ipni.org:names:77209990-1

Figures 1, 2A–F, 3, 4A–C, 5

Type. CHINA. Guizhou: Weining County, Jinzhong Town, 2226 m alt., 5 Aug 2018, X.X. Zhu et al. ZXX18217 (holotype: CSH–0172289!; isotypes: CSH!, KUN!).

Diagnosis. *Isotrema brevilimbus* is morphologically similar to *I. ovatifolium* (S.M. Hwang) X.X. Zhu, S. Liao & J.S. Ma and *I. wardianum* (J.S. Ma) X.X. Zhu, S. Liao




Figure 1. *Isorema brevilimbium* X.X.Zhu, Jun Wang & F.Cao. **A** Branch **B** leaf **C, D** flower **E** longitudinal-section of flower (showing inside structure) **F** anthers and gynostemium **G** capsule **H** seed. Drawn by S.Z. Qiao.




Figure 2. **A–F** *Isotrema brevilibum* X.X.Zhu, Jun Wang & F.Cao. **A** Habitat **B** leaves **C** lateral view of flower **D** frontal view of flower **E** anthers and gynostemium **F** capsule **G–I** *I. wardianum* **G** habit **H** frontal view of flower **I** anthers and gynostemium. **A** Photographed by F. Cao **B, E, F** photographed by X.X. Zhu **C, D** photographed by G. Liu **G** photographed by C. Liu **H, I** photographed by J.D. Ya.

& J.S. Ma, but differs from the former in its lamina long ovate (vs. lamina ovate in *I. ovatifolium*), perianth limb forming right angle with upper tube, length nearly equal to width, and apex dark purple and opened (vs. limb straightly extended from upper tube, length significantly longer than width, and apex dark purple and constricted in *I. ovatifolium*), differs from the latter in its lamina long ovate and abaxially densely villous (vs. lamina lanceolate and abaxially subglabrous or glabrous in *I. wardianum*), perianth limb forming right angle with upper tube, length nearly equal to width, and apex dark purple and opened (vs. limb forming obtuse angle with upper tube, length significantly longer than width, and apex light yellow and constricted in *I. wardianum*).



Figure 3. Holotype of *Isotrema brevilibum* X.X.Zhu, Jun Wang & F.Cao (CSH-0172289).




Figure 4. Leaves, lateral view of flowers, and longitudinal dissected flowers of *Isotrema brevilimbium* (A–C), *I. ovatifolium* (D–F), and *I. wardianum* (G–I). A–F Photographed by X.X. Zhu G photographed by C. Liu H, I photographed by J.D. Ya.

Description. Climbing shrubs. Stems terete, densely villous when young, old branchlets glabrous. Petioles 1–4 cm long, densely villous; laminas long ovate, 5–13 × 2.5–3.5 cm, adaxially appressed villous, abaxially densely villous, base cordate, margin entire, apex acute; basal veins palmate, 2–3 pairs from base, lateral veins 4–6-paired. Flowers axillary or lateral on young stems, solitary, rarely paired. Pedicels pendulous, 1.5–3 cm long, densely villous; bracteole ovate, conduplicate, ca. 2 × 1 mm, abaxially densely villous, adaxially smooth, inserted on lower part of pedicel. Perianth tube geniculately curved, abaxially villous; basal tube ca. 1 cm long, inside dark red, upper tube ca. 1.5 cm long, inside red; limb short cylinder, length nearly equal to width, ca. 7 × 8 mm, forming right angle with upper tube, apex dark purple, opened, ca. 7 mm wide at mouth, inside dark red with densely tiny dark-purple papillae; throat subcircular, ca. 4 mm wide. Anthers 6, oblong, ca. 1.5 mm




Figure 5. Distribution of *Isotrema brevilibum*, *I. ovatifolium*, and *I. wardianum* based on field observation, specimens and literatures examined.

long, adnate in 3 pairs to base of gynostemium, opposite to lobes. Gynostemium ca. 3 mm long, 3-lobed. Ovary terete, ca. 8 mm long, densely villous. Capsule cylindric, abaxially densely villous, ca. 4.5 × 2 cm. Seeds ovate, 4–5 × 3–3.5 mm, concave-convex.

Phenology. Flowering from May to August, fruiting from July to September.

Etymology. The specific epithet refers to the short cylinder perianth limb of the new species. The “brevi” means “short”, “limbum” means “limb”, so the new species is named *Isotrema brevilibum*.

Common name (assigned here). Duan Yan Guan Mu Tong (短檐关木通; Chinese name).

Distribution and habitat. The new species is currently only known from Weining County of Guizhou, China. It grows by the roadside of farmland at an altitude of ca. 2200 m.

Preliminary conservation status. *Isotrema brevilibum* is known from a single population including two individuals on the roadside of farmland. The new species is assigned a preliminary status of vulnerable (VU) according to the IUCN Red List Categories (IUCN 2012). However, since very few details exist about its natural distribution, the lack of sufficient data currently does not allow a final risk evaluation and the species might be regarded as data deficient (DD). Further field surveys in western Guizhou and northeastern Yunnan are needed to gain more information on its distri-

bution. Not only is the area not under protection as a nature reserve, but also habitat disturbance brought about by human activities, such as grazing and farming, may have a negative impact on the new species.

Note. *Isotrema wardianum* was previously only known from Myanmar and India. Sun and Zhou (2002) later reported the species from China, according to a specimen collected from Medog County of Tibet (*H. Sun et al.* 4935), but without flower or fruit. Nevertheless, the species had long been neglected by taxonomic studies of Huang et al. (2003), Do et al. (2015a), and Zhu et al. (2019a, 2019d) on Chinese *Isotrema*. It was not until 2018 that we discovered a seedling of *Isotrema* sp. at the same locality as that of *H. Sun et al.* 4935 and transplanted it in the nursery of the Kunming Institute of Botany. A year later, the plant grown from this seedling bloomed and enabled us to identify it as *I. wardianum* (Figs 2G–I, 4G–I) and confirm its distribution in China.

Discussion

Isotrema brevilimbium is morphologically similar to *I. ovatifolium* and *I. wardianum* in the shape, size, and color of flower and the dark-purple papillae in the inner surface of perianth limb, but they can be distinguished by the morphology of lamina, the angle between perianth limb and upper tube, as well as the length and mouth of limb. Detailed morphological comparisons among the three species are summarized in Table 1 and Fig. 4.

Specimens of *Isotrema wardianum* examined. MYANMAR. Adung Valley, 12 Apr 1931, F. Kingdon-Ward 9398 (holotype: BM). CHINA. Tibet: Medog County, 2100 m, 21 Mar 1993, H. Sun et al. 4935 (KUN); **at the same locality**, 1705 m, 27 Nov 2018, C. Liu & J.D. Ya 18CS17145 (KUN).

Specimens of *Isotrema ovatifolium* examined. CHINA. Guizhou: Weining County, Jinzhong Town, 2232 m, 5 Aug 2018, Zhu et al. ZXX18218 (CSH, KUN); Sichuan: Huidong County, 2520 m, 27 Jun 1959, S.K. Wu 1584 (type: SM).

Table 1. Morphological comparisons among *Isotrema brevilimbium*, *I. ovatifolium* and *I. wardianum*. These characters were based on field observation, related specimens and literatures (Hwang 1981; Ma 1989a; Huang et al. 2003).

Characters	<i>I. brevilimbium</i>	<i>I. ovatifolium</i>	<i>I. wardianum</i>
Lamina	long ovate, 5–13 × 2.5–3.5 cm, abaxially densely villous, base cordate	ovate, 5–13 × 4–8 cm, abaxially densely villous, base cordate	lanceolate, 12–16 × 3–4 cm, abaxially subglabrous or glabrous, base auriculate
Perianth limb	short cylinder, forming right angle with upper tube, length nearly equal to width, apex dark purple, opened, ca. 7 mm wide at mouth	cylinder, straightly extended from upper tube, length significantly longer than width, apex dark purple, constricted, ca. 1 mm wide at mouth	cylinder, forming obtuse angle with upper tube, length significantly longer than width, apex light yellow, constricted, ca. 3 mm wide at mouth
Perianth throat	ca. 4 mm wide	ca. 2.5 mm wide	ca. 2 mm wide
Anthers	ca. 1.5 mm long	ca. 1.5 mm long	ca. 2 mm long
Gynostemium	ca. 3 mm long	ca. 3.5 mm long	ca. 3.5 mm long
Capsule	ca. 4.5 × 2 cm	ca. 6 × 2 cm	unknown
Distribution	China (Guizhou)	China (Guizhou, Sichuan, Yunnan)	China, Myanmar, India

Acknowledgements

The authors are very grateful to Ms. Shi-Zhen Qiao for the line drawing of *I. brevilibum*. This study was jointly supported by the National Natural Science Foundation of China (grant nos. 31600161 and 31370225), the Nanhu Scholars Program for Young Scholars of XYNU, and the Institute for Conservation and Utilization of Agro-bioresources in the Dabie Mountains.

References

- Cai L, He DM, Huang YS, Dao ZL (2020a) *Aristolochia wenshanensis*, a new species of Aristolochiaceae from karst region in southeastern Yunnan, China. *Taiwania* 65(1): 41–46.
- Cai L, Dao ZL, Zhu XX (2020b) *Isotrema hei* (Aristolochiaceae), a new species from southeastern Yunnan, China. *Annales Botanici Fennici* 57: 125–129. <https://doi.org/10.5735/085.057.0117>
- Do TV, Nghiem TD (2017) Taxonomic notes on some *Aristolochia* species in Vietnam. *Taiwania* 62(2): 216–218.
- Do TV, Luu TH, Wanke S, Neinhuis C (2015a) Three new species and three new records of *Aristolochia* subgenus *Siphisia* from Vietnam including a key to the Asian species. *Systematic Botany* 40(3): 671–691. <https://doi.org/10.1600/036364415X689140>
- Do TV, Neinhuis C, Wanke S (2015b) A new species of *Aristolochia* subgenus *Siphisia* (Aristolochiaceae) from central Vietnam. *Phytotaxa* 220(1): 69–76. <https://doi.org/10.11646/phytotaxa.220.1.6>
- Do TV, Nghiem TD, Wanke S, Neinhuis C (2014) *Aristolochia quangbinhensis* (Aristolochiaceae), a new species from Central Vietnam. *PhytoKeys* 33: 51–59. <https://doi.org/10.3897/phytokeys.33.6094>
- Do TV, Nguyen DQ, Nguyen TQT, Wanke S, Neinhuis C (2015c) *Aristolochia cochinchinensis* (Aristolochiaceae), a new species from southern Vietnam. *Annales Botanici Fennici* 52(3–4): 268–273. <https://doi.org/10.5735/085.052.0321>
- Do TV, Truong CQ, Huynh HTT (2017) *Aristolochia neinhuisii* (Aristolochiaceae), a new species from Vietnam. *Annales Botanici Fennici* 54(4–6): 203–208. <https://doi.org/10.5735/085.054.0602>
- Do TV, Vu TTH, Luu HT, Nguyen TT (2019) *Aristolochia nuichuaensis* (subg. *Siphisia*, Aristolochiaceae), a New Species, an updated key and a checklist to the species of *Siphisia* in Vietnam. *Annales Botanici Fennici* 56(1–3): 107–113. <https://doi.org/10.5735/085.056.0116>
- Do TV, Wanke S, Neinhuis C (2016) *Aristolochia bidoupensis* sp. nov. from southern Vietnam. *Nordic Journal of Botany* 34(5): 513–516. <https://doi.org/10.1111/njb.01066>
- Do TV, Wanke S, Neinhuis C, Pooma R (2015d) *Aristolochia phuphathanaphongiana* sp. nov. from southwestern Thailand. *Nordic Journal of Botany* 33(5): 567–571. <https://doi.org/10.1111/njb.00889>
- Gong QB, Landrein S, Xi HC, Ma XD, Yang ZH, He KW, Shen JY (2018) *Aristolochia tongbi-guanensis*, a new species of Aristolochiaceae from Yunnan, China. *Taiwania* 63(3): 183–187.
- Hwang SM (1981) Materials for Chinese *Aristolochia*. *Zhiwu Fenlei Xuebao* 19: 222–231.

- Huang SM, Kelly LM, Gilbert MG (2003) *Aristolochia* Linnaeus. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* 5. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 258–269.
- Huang YS, Peng RC, Tan WN, Wei GF, Liu Y (2013) *Aristolochia mulunensis* (Aristolochiaceae), a new species from limestone areas in Guangxi, China. *Annales Botanici Fennici* 50(3): 175–178. <https://doi.org/10.5735/085.050.0308>
- Huang YS, Peng YD, Huang BY, Lv HZ, Lin CR (2015) *Aristolochia gongchengensis* (Aristolochiaceae), a new species from the limestone areas in Guangxi, China. *Annales Botanici Fennici* 52(5–6): 396–400. <https://doi.org/10.5735/085.052.0522>
- Hwang SM (1988) *Aristolochia* Linnaeus. In: Kiu HS, Ling YR (Eds) *Flora Reipublicae Popularis Sinicae* 24. Science Press, Beijing, 199–245.
- Huong NTT, Hai DV, Quang BH, Cuong NT, Khang NS, Vu DQ, Ma JS (2014) *Aristolochia xuanlienensis*, a new species of Aristolochiaceae from Vietnam. *Phytotaxa* 188(3): 176–180. <https://doi.org/10.11646/phytotaxa.188.3.7>
- IUCN (2012) IUCN Red List Categories and Criteria, Version 3.1 (2nd Ed.) Gland and Cambridge, 32 pp.
- Li RT, Wang ZW, Wang J, Zhu XX, Xu H (2019) *Isotrema sanyaense*, a new species of Aristolochiaceae from Hainan, China. *PhytoKeys* 128: 85–96. <https://doi.org/10.3897/phytokeys.128.35042>
- Liu ZW, Deng YF (2009) *Aristolochia wuana*, a new name in Chinese *Aristolochia* (Aristolochiaceae). *Novon* 19(3): 370–371. <https://doi.org/10.3417/2007151>
- Lu CT, Wang JC (2014) *Aristolochia yujungiana* (Aristolochiaceae): A new species from Taiwan. *Taiwan Linye Kexue* 29: 291–299.
- Ma JS (1989a) A revision of *Aristolochia* Linn. from E. & S. Asia. *Zhiwu Fenlei Xuebao* 27: 321–364.
- Ma JS (1989b) A revision of *Aristolochia* from Yunnan. *Yunnan Zhi Wu Yan Jiu* 11: 321–323.
- Ohi-Toma T, Watanabe-Toma K, Murata H, Murata J (2014) Morphological variations of *Aristolochia kaempferi* and *A. tanzawana* (Aristolochiaceae) in Japan. *Shokubutsu Kenkyu Zasshi* 89: 152–163.
- Peng YD, Gadagkar SR, Li J, Xie YY, Huang XY, Lu HZ, Huang BY, Yu LY (2019) *Aristolochia kechangensis* sp. nov (Aristolochiaceae) from Guangxi, China. *Nordic Journal of Botany* 37(9): e02456. <https://doi.org/10.1111/njb.02456>
- Sun H, Zhou ZK (2002) *Seed Plants of The Big Bend Gorge of Yalu Tsangpo in SE Tibet, E Himalayas*. Yunnan Science & Technology Press, Kunming, 425 pp.
- Tao DD (1997) *Aristolochia* Linnaeus. In: Chen SK (Ed.) *Flora Yunnanica* 8. Science Press, Beijing, 7–25.
- Thiers B (2020) *Index Herbariorum: A global directory of public herbaria and associated staff*. New York Botanical Garden's virtual Herbarium. <http://sweetgum.nybg.org/ih/> [accessed 11 March 2020]
- Wu L, Xu WB, Huang YS, Liu Y (2015) *Aristolochia longlinensis* (Aristolochiaceae), a new species from Western Guangxi, China. *Novon* 23(4): 490–493. <https://doi.org/10.3417/2011105>
- Wu L, Xu WB, Wei GF, Liu Y (2013) *Aristolochia huanjiangensis* (Aristolochiaceae), a new species from Guangxi, China. *Annales Botanici Fennici* 50(6): 413–416. <https://doi.org/10.5735/085.050.0608>

- Xu H, Li YD, Yang HJ, Chen HQ (2011) Two new species of *Aristolochia* (Aristolochiaceae) from Hainan Island, China. *Novon* 21(2): 285–289. <https://doi.org/10.3417/2009116>
- Yang B, Ding HB, Zhou SS, Zhu XX, Li R, Mya BM, Tan YH (2018) *Aristolochia sinoburmanica* (Aristolochiaceae), a new species from north Myanmar. *PhytoKeys* 94: 13–22. <https://doi.org/10.3897/phytokeys.94.21557>
- Yao TL (2012) *Aristolochia vallisicola* (Aristolochiaceae), a new species from Peninsular Malaysia. *PhytoKeys* 14: 15–22. <https://doi.org/10.3897/phytokeys.14.3354>
- Zhou XX, Jiang GB, Zhu XX, Liu ZY, Huang Y, Wang GT, Wang RJ (2019) *Isotrema plagiostomum* (Aristolochiaceae), a new species from Guangdong, South China. *Phytotaxa* 405(4): 221–225. <https://doi.org/10.11646/phytotaxa.405.4.7>
- Zhu XX, Li XQ, Liao S, Du C, Wang Y, Wang ZH, Yan J, Zuo YJ, Ma JS (2019a) Reinstatement of *Isotrema*, a new generic delimitation of *Aristolochia* subgen. *Siphisia* (Aristolochiaceae). *Phytotaxa* 401(1): 1–23. <https://doi.org/10.11646/phytotaxa.401.1.1>
- Zhu XX, Li XQ, Liao S, Li GD, Ma JS (2019b) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) V: Two new species from Yunnan, China. *PhytoKeys* 130: 93–106. <https://doi.org/10.3897/phytokeys.130.33933>
- Zhu XX, Liao S, Zhang L, Wang ZH, Du C, Ma JS (2016) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) I: Confirmation and illustration of *A. austroszechuanica* *A. faucimaculata* and *A. yunnanensis* var. *meionantha* from China. *Phytotaxa* 261(2): 137–146. <https://doi.org/10.11646/phytotaxa.261.2.3>
- Zhu XX, Liao S, Sun ZP, Zhen AG, Ma JS (2017a) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) II: Identities of *Aristolochia austroyunnanensis* *A. dabieshanensis* and *A. hyperxantha* – a new species from Zhejiang, China. *Phytotaxa* 313(1): 61–76. <https://doi.org/10.11646/phytotaxa.313.1.4>
- Zhu XX, Liao S, Ma ZX, Xu B, Wang ZH, Wang Y, Ma JS (2017b) The taxonomic revision of Asian *Aristolochia* (Aristolochiaceae) III: Two new taxa of *Aristolochia* and morphological revision for the flower character of *A. obliqua* from Yunnan, China. *Phytotaxa* 332(3): 269–279. <https://doi.org/10.11646/phytotaxa.332.3.3>
- Zhu XX, Shen B, Sun ZP, Chen B, Liao S, Ma JS (2018) Two New Species of *Aristolochia* (Aristolochiaceae) from Yunnan, China. *Novon* 26(3): 298–306. <https://doi.org/10.3417/2018066>
- Zhu XX, Wang J, Liao S, Ma JS (2019d) Synopsis of *Aristolochia* L. and *Isotrema* Raf (Aristolochiaceae) in China. *Shengwu Duoyangxing* 27(10): 1143–1146. <https://doi.org/10.17520/biods.2019183>
- Zhu XX, Zhang L, Hua ZX, Chen GF, Liao S, Ma JS (2015) *Aristolochia weixiensis*, a new species of Aristolochiaceae from Yunnan, China. *Phytotaxa* 230(1): 54–60. <https://doi.org/10.11646/phytotaxa.230.1.4>
- Zhu XX, Zheng HL, Wang J, Gao YQ, Ma JS (2019c) Taxonomic studies on the genus *Isotrema* (Aristolochiaceae) from China: I. *I. cangshanense*, a new species from Yunnan. *PhytoKeys* 134: 115–124. <https://doi.org/10.3897/phytokeys.134.37243>