

Hoya spectatissima (Gentianales, Apocynaceae), a new species from Yunnan, China

Bine Xue¹, Erfeng Huang², Gang Yao³, Guohua Zhao⁴, Jiuxiang Huang³

1 College of Horticulture and Landscape Architecture, Zhongkai University of Agriculture and Engineering, Guangzhou 510225, China **2** Guangxi Nanning Roy Garden Co., Ltd., Nanning 530227, China **3** College of Forestry and Landscape Architecture, South China Agricultural University, Guangzhou 510642, China **4** Fairy Lake Botanical Garden, Shenzhen & Chinese Academy of Sciences, Shenzhen 518004, Guangdong, China

Corresponding author: Jiuxiang Huang (jxhuang@scau.edu.cn)

Academic editor: Petra De Block | Received 30 December 2022 | Accepted 2 March 2023 | Published 21 March 2023

Citation: Xue B, Huang E, Yao G, Zhao G, Huang J (2023) *Hoya spectatissima* (Gentianales, Apocynaceae), a new species from Yunnan, China. *PhytoKeys* 222: 69–74. <https://doi.org/10.3897/phytokeys.222.99420>

Abstract

Hoya spectatissima, a new species from Yunnan Province, China, is described and illustrated. *Hoya spectatissima* is morphologically similar to *H. lyi*, but can be easily distinguished from the latter by its succulent leaves that are 2–4.5 cm long (vs. coriaceous leaves that are up to 9 cm long), corolla that is pink to pale pink (vs. yellowish-white) and corona lobes that are sub-rhombic in top view (vs. ovoid in top view).

Keywords

Asclepiadoideae, Asia, Marsdenieae, morphology, nomenclature, taxonomy

Introduction

Hoya R. Br. is a large genus within the tribe Marsdenieae of Apocynaceae (Wanntorp et al. 2014), comprising 350–450 species of subshrubs or lianas widely distributed in the tropical and subtropical regions of Asia, Oceania and the Pacific Islands (Li et al. 1995; Endress et al. 2018). China possesses more than 40 species (Huang et al. 2021; Huang et al. 2022), with multiple new species or newly-recorded species reported recently, such as *H. yingjiangensis* J.Feng Zhang, L.Bai, N.H.Xia & Z.Q.Peng (Zhang et al. 2015), *H. acuminata* (Wight) Benth. ex Hook.f. (Gui et al. 2017), *H. vangviengensis* Rodda & Simonsson (Zhang et al. 2017), *H. tamdaoensis* Rodda & T.B.Tran (Nong et

al. 2018), *H. burmanica* Rolfe (Ma et al. 2019), *H. longicalyx* Wang Hui & E.F.Huang (Huang et al. 2020), *H. gaoligongensis* M.X.Zhao & Y.H.Tan (Zhao et al. 2020), *H. ny-
ingchiensis* Y.W.Zuo & H.P.Deng (Zuo et al. 2020), *H. pyrifolia* E.F. Huang (Huang
et al. 2021), *H. longlingensis* E.F. Huang and *H. sichuanensis* E.F.Huang (Huang et al.
2022). Recently, we found an unknown *Hoya* species (Fig. 1) in southwest Yunnan
Province, China. After a detailed morphological comparison with all the *Hoya* species
recorded in China and adjacent regions, we concluded that this species is new to sci-
ence. Thus, we formally describe it here.

Taxonomic treatment

Hoya spectatissima B.Xue, E.F.Huang, Gang Yao & J.X.Huang, sp. nov.

urn:lsid:ipni.org:names:77316067-1

Fig. 1

Diagnosis. *Hoya spectatissima* is most similar to *H. lyi* H.Lév, but differs from the lat-
ter by its succulent leaves (vs. coriaceous leaves) that are 2–4.5 cm in length (vs. up
to 9 cm in length) and rounded or rarely obtuse at the apex (vs. mostly acute at the
apex), pink to pale pink corolla (vs. yellowish-white corolla) that is densely sericeous
adaxially (vs. densely pubescent adaxially), ivory white corona lobes (vs. purple towards
the inner angle and ivory white towards the outer angle of corona lobes) that are sub-
rhombic in top view (vs. ovoid in top view) and have an acute outer angle (vs. truncate
to shallowly concave outer angle).

Type. CHINA, Yunnan Province, Gengma Hsien, Sipaishan Town, Guanlong Vil-
lage, Luoshuidong, on rocks in karst habitat, 7 July 2021, *E.F. Huang-RG0017* (Holo-
type: IBSC!).

Description. Lithophytic climber. Stems rounded in cross section, up to 1 m in
length, 1.5–2 mm in diameter, densely pubescent when young, subglabrous when
mature, sometimes 3–5 branches at nodes, internodes 3–8 cm long. Leaves opposite,
wine-red and pubescent when young; mature lamina subglabrous, succulent, elliptic or
oblong, 2–4.5 × 1.3–1.8 cm, 2.5–3 mm in thickness, bright green adaxially, pale green
abaxially, base rounded, apex rounded or rarely obtuse; mid-vein slightly elevated or
obscure adaxially, not elevated abaxially, lateral veins 2–3 pairs or sometimes invisible
in adaxial surface, invisible in abaxial surface; petioles sometimes recurved, 3–7 mm
long, 1–1.5 mm in diameter. Inflorescences axillary pseudo-umbels, many-flowered,
globose; peduncle longer than pedicels, 4.5–5 cm long, papillose to sparsely pubescent;
pedicels 2.5–3 cm long, yellow-green, papillose to sparsely pubescent. Calyx lobes tri-
angular to narrowly triangular, ca. 1.5 × 2 mm, purplish-red, sparsely pubescent or
glabrous. Corolla rotate, pink in early flowering phase, pale pink in late flowering
phase, ca. 2 cm in diameter, densely sericeous adaxially, sub-glabrous abaxially, lobes




Figure 1. *Hoya spectatissima* **A** habit **B** cross section of lamina **C** leaves **D, E** inflorescence **F** flower (top view) **G** flower (bottom view) **H** corona (top view; Red arrowheads indicate the inner angle of the corona lobes and yellow arrowheads indicate the outer angle of the corona lobes) **I** corona (side view) **J** ovaries **K** pollinarium.




Figure 2. *Hoya luyi* **A** isosyntype (*E. Bodinier* 1853, P00634485) **B** inflorescence **C** habit.

triangular, 5–6 mm long and 5–6 mm wide, margin recurved, apex rounded, revolute. Corona ivory white, 7–8 mm in diameter, 2.0–2.4 mm high, lobes 5, stellate spreading, ca. 3.5×2.5 mm, sub-rhombic in top view, outer angle acute, inner angle acuminate, the inner tips convex and spreading obviously towards the centre of the flower, ivory white or rarely shallowly pink. Pollinia clavate, ca. 0.73×0.26 mm, base cuneate, apex truncate, narrowing towards the base, sterile edge all along the outer edge of the pollinium, translator arms attached at the centre of the corpusculum. Ovaries 2, triangular-ovate, attached to each other, ca. 2 mm long and 1.5 mm wide, green, glabrous. Fruit and seed not seen.

Distribution and habit. *Hoya spectatissima* is known from at least two localities in Cangyuan Hsien and Gengma Hsien, Yunnan Province, China. It is a lithophytic liana that climbs on rocks in karst habitat. It is a common species in Guanlong Village, Gengma Hsien, but an occasional species in Menglai Town, Cangyuan Hsien.

Etymology. *Hoya spectatissima* is named to reflect its beautiful flowers.

Taxonomic discussion. *Hoya spectatissima* has evidently succulent leaves that are 2.5–3 mm thick (Fig. 1B); this trait is also found in *H. pandurata* Tsiang, which is also endemic in Yunnan Province, China. However, *H. pandurata* is a subshrub and is epiphytic on trees in open or mixed woods and it further differs from *H. spectatissima* by its pandurate or oblong leaves (vs. elliptic or oblong) with acuminate apex (vs. rounded or rarely obtuse), shorter peduncle and pedicels that are ca. 3 mm long (vs.

4.5–5 cm long) and 1.5 cm long (vs. 2.5–3 cm long) respectively and yellow or reddish corolla (vs. pink or pale pink) that is 0.8–1 cm (vs. ca. 2 cm) in diameter (Li et al. 1995). Additionally, *H. spectatissima* is morphologically similar to *H. lyi*, a species recorded from China, Laos and Vietnam (Rodda 2012). Both species have elliptic or oblong leaves (Figs 1A, C, 2A, C), triangular to narrowly triangular calyx lobes, corolla ca. 2 cm in diameter and clavate pollinia. However, *Hoya spectatissima* differs from *H. lyi* by the following characters: succulent leaves that are 2.5–3 mm thick (Fig. 1B) (vs. coriaceous leaves that are less than 1.5 mm thick, Fig. 2A, C); shorter leaves that are 2–4.5 cm long (Fig. 1C) (vs. up to 9 cm long, Fig. 2A); apex of leaves rounded or rarely obtuse (Fig. 1A, C) (vs. mostly acute, Fig. 2A, C); 2–3 paired lateral veins in leaves (vs. 3–5 paired lateral veins); pink to pale pink corolla (Fig. 1D, E) (vs. yellowish-white, Fig. 2B); corolla lobes densely sericeous adaxially (Fig. 1F, G) (vs. densely pubescent adaxially, Fig. 2B); corona lobes sub-rhombic in top view (Fig. 1F, H) (vs. ovoid, Fig. 2B) and ivory white (Fig. 1F, H) (vs. purple towards the inner angle and ivory white towards the outer angle, Fig. 2B); and outer angle of corona lobes acute (Fig. 1H) (vs. truncate to shallowly concave, Fig. 2B).

Paratype. China. Yunnan Province, Cangyuan Hsien, Menglai Town, on rocks in karst habitat, 22 August 2021, *E.F. Huang-RG0033* (IBSC).

Acknowledgements

The authors are indebted to the curators of the Herbaria E, IBSC, K, KUN, NY, P and PE for providing specimen images or hosting our visits, to L.Y. Li in the Kunming Institute of Botany, Chinese Academy of Sciences, for the photograph in Fig. 1K. This study was financially supported by the National Science Foundation of China (Grant no. 31872646).

References

- Endress ME, Meve U, Middleton DJ, Schumann SL (2018) Apocynaceae. In: Kadereit JW, Bittrich V (Eds) Flowering Plants. Eudicots, The Families and Genera of Vascular Plants (Vol. 15). Springer, Cham, Switzerland, 207–411. https://doi.org/10.1007/978-3-319-93605-5_3
- Gui LJ, Peng YD, Huang BY, Lan XZ, Yu LY (2017) *Hoya acuminata* (Wight) Benth. ex Hook. f. (Apocynaceae, Asclepiadoideae), a newly recorded species from China. *Acta Botanica Boreali-Occidentalia Sinica* 37: 1252–1254. <https://doi.org/10.7606/j.issn.1000-4025.2017.06.1252>
- Huang EF, Yang LL, Xi W, Liu YY, Wang H (2020) *Hoya longicalyx*, a new species of *Hoya* (Apocynaceae: Asclepiadoideae) from Yunnan, China. *Taiwania* 65: 353–359.
- Huang EF, Yao G, Jiang RH, Yang LL, Xi W, Zhang ZS, Zhang XC (2021) *Hoya pyriformis* (Apocynaceae), a new species from south-western Yunnan, China. *PhytoKeys* 174: 95–106. <https://doi.org/10.3897/phytokeys.174.60137>

- Huang EF, Tang LM, Zhang ZF, Huang JX, Ou ZF, Zhang XC (2022) *Hoya longlingensis* and *H. sichuanensis* (Apocynaceae), two new species from Southwestern China. *Taxonomy* 2(1): 99–106. <https://doi.org/10.3390/taxonomy2010008>
- Li PT, Gilbert MG, Stevens WD (1995) *Hoya*. In: Wu CY, Raven PH (Eds) *Flora of China* (Vol. 16). Science Press & Missouri Botanical Garden Press, Beijing, China / St. Louis, 228–236.
- Ma X, Gong Q, Yang Z, Shi J, Shen J (2019) *Hoya burmanica*, a newly recorded species of *Hoya* (Asclepiadaceae) from China. *Acta Botanica Boreali-Occidentalia Sinica* 39(5): 948–949. <https://doi.org/10.7606/j.issn.1000-4025.2019.05.0948>
- Nong DX, Peng YD, Huang BY, Yu LY (2018) *Hoya tamdaoensis*, a newly recorded species of Apocynaceae from China. *Redai Yaredai Zhiwu Xuebao* 26(1): 82–84.
- Rodda M (2012) Taxonomy of *Hoya lyi*, *Hoya yuennanensis* and *Hoya mekongensis* (Apocynaceae – Asclepiadoideae). *Edinburgh Journal of Botany* 69(1): 83–93. <https://doi.org/10.1017/S0960428611000412>
- Wanntorp L, Grudinski M, Forster PI, Muellner-Riehl A, Grimm GW (2014) Wax plants (*Hoya*, Apocynaceae) evolution: Epiphytism drives successful radiation. *Taxon* 63(1): 89–102. <https://doi.org/10.12705/631.3>
- Zhang JF, Bai L, Xia N, Peng Z (2015) *Hoya yingjiangensis* (Apocynaceae, Asclepiadoideae), a new campanulate-flowered species from Yunnan, China. *Phytotaxa* 219(3): 283–288. <https://doi.org/10.11646/phytotaxa.219.3.8>
- Zhang JF, Xia NH, Li DM, Li S (2017) *Hoya vangviengensis*, a newly recorded species of *Hoya* (Asclepiadoideae, Apocynaceae) from China. *Redai Yaredai Zhiwu Xuebao* 25(6): 594–596.
- Zhao MX, Wang HY, Yin ZJ, Wang MJ, Yang JC, Tan YH (2020) *Hoya gaoligongensis* (Apocynaceae, Asclepiadoideae), a new species from Yunnan, SW China. *Phytotaxa* 459(3): 219–226. <https://doi.org/10.11646/phytotaxa.459.3.3>
- Zuo YW, Lan XZ, Deng HP, Zhao FY, Yuan XF (2020) *Hoya nyingchiensis* (Apocynaceae, Asclepiadoideae), a new species from Tibet, China. *Phytotaxa* 468(1): 130–136. <https://doi.org/10.11646/phytotaxa.468.1.9>